


Chapter 22. Fatty Acid Metabolism

Chem 454: Biochemistry II
University of Wisconsin-Eau Claire




Introduction

Fatty acids play several important roles:


- Building blocks for phospholipids and glycolipids
- Target proteins to membranes
- High energy source of fuel
- Fatty acid derivatives are used as hormones and intracellular messengers

Overview of fatty acid synthesis




Introduction

FATTY ACID DEGRADATION




FATTY ACID SYNTHESIS



1. Triglycerides


Triglycerides are a highly concentrated store of energy

- 9 kcal/g vs 4 kcal/g for glycogen
- Glycogen is also highly hydrated, 2 g H₂O/g glycogen




1.1 Pancreatic Lipases

- Dietary triacylglycerols must be broken down before being absorbed by the intestines.
- Bile salts, which act as detergents, are used to solublize the triacylglycerols




1.1 Pancreatic Lipases

- Dietary triacylglycerols must be broken down before being absorbed by the intestines.
- Bile salts, which act as detergents, are used to solublize the triacylglycerols




1.1 Pancreatic Lipases

Pancreatic lipases hydrolyze the ester bonds of the triacylglycerols while in the micelles.



1.1 Chylomicrons

In the intestinal mucosal cells, the fatty acids and monoacylglycerides are resynthesized into triacylglycerides and packaged into *chylomicrons*. Chylomicrons and lymph are dumped via the thoracic duct into the left subclavian vein




1.1 Chylomicrons

Chylomicrons and lymph are dumped via the thoracic duct into the left subclavian vein.

Want to know more about lymphatic system?

Try here:

<http://owensboro.kctcs.edu/gcaplan/ant2/notes/Notes7%20Lymphatic%20Anatomy.htm>



Copyright © 2001 Benjamin Cummings, an imprint of Addison Wesley Longman, Inc.


2. Utilization of Fatty Acids as Fuel

Three stages of processing

- Triglycerols are degraded to fatty acids and glycerol in the adipose tissue and transported to other tissues.
- Fatty acids are activated and transported into the mitochondria.
- Fatty acids are broken down into two-carbon acetyl-CoA units and fed into the citric acid cycle.

2.1 Breakdown of Triacylglycerols

In the adipose tissue, lipases are activated by hormone signaled phosphorylation



2.1 Breakdown of Triacylglycerols





Figure 22-7
Biochemistry, Sixth Edition
© 2007 W. H. Freeman and Company

2.1 Breakdown of Triacylglycerols


The lipases break the triacylglycerols down to fatty acids and glycerol

- The fatty acids are transported in the blood by serum albumin





2.1 Breakdown of Triacylglycerols

The glycerol is absorbed by the liver and converted to glycolytic intermediates.




2.2 Activation of Fatty Acids

Acyl CoA synthetase reaction occurs in the on the mitochondrial membrane.




2.3 Transport into Mitochondrial Matrix

Carnitine carries long-chain activated fatty acids into the mitochondrial matrix




2.3 Transport into Mitochondrial Matrix

Carnitine carries long-chain activated fatty acids into the mitochondrial matrix



2.3 Transport into Mitochondrial Matrix


A Miracle???



2.4 Fatty acid oxidation

Each round in fatty acid degradation involves four reactions


- 1. oxidation to
- *trans*- Δ^2 -Enoyl-CoA



2.4 Fatty acid oxidation

Each round in fatty acid degradation involves four reactions


- 2. Hydration to L-3-Hydroxyacyl CoA



2.4 Fatty acid oxidation

Each round in fatty acid degradation involves four reactions


- 3. Oxidation to
- 3-Ketoacyl CoA



2.4 Fatty acid oxidation

Each round in fatty acid degradation involves four reactions


- 4. Thiolytic cleavage to produce Acetyl-CoA



2.4 Fatty acid oxidation

Each round in fatty acid degradation involves four reactions

- The process repeats itself



2.4 Fatty acid oxidation


Each round in fatty acid degradation involves four reactions

TABLE 22.1 Principal reactions in fatty acid oxidation

Step	Reaction	Enzyme
1	Fatty acid + CoA + ATP \rightleftharpoons acyl CoA + AMP + PP _i	Acyl CoA synthetase [also called fatty acid thiokinase and fatty acid:CoA ligase (AMP)]
2	Carnitine + acyl CoA \rightleftharpoons acyl carnitine + CoA	Carnitine acyltransferase (also called carnitine palmitoyl transferase)
3	Acyl CoA + E-FAD \rightarrow <i>trans</i> - Δ^2 -enoyl CoA + E-FADH ₂	Acyl CoA dehydrogenases (several isozymes having different chain-length specificity)
4	<i>trans</i> - Δ^2 -Enoyl CoA + H ₂ O \rightleftharpoons L-3-hydroxyacyl CoA	Enoyl CoA hydratase (also called crotonase or 3-hydroxyacyl CoA hydrolyase)
5	L-3-Hydroxyacyl CoA + NAD ⁺ \rightleftharpoons 3-ketoacyl CoA + NADH + H ⁺	L-3-Hydroxyacyl CoA dehydrogenase
6	3-Ketoacyl CoA + CoA \rightleftharpoons acetyl CoA + acyl CoA (shortened by C ₂)	β -Ketothiolase (also called thiolase)


2.5 ATP Yield

The complete oxidation of the sixteen carbon palmitoyl-CoA produces 106 ATP's



3.1 Special Cases

Unsaturated fatty acids (monounsaturated)



3.1 Special Cases

Unsaturated fatty acids (polyunsaturated)




Figure 22-11
Biochemistry, Sixth Edition
© 2007 W.H. Freeman and Company

3.2 Odd-Chain





Figure 22-12
Biochemistry, Sixth Edition
© 2007 W. H. Freeman and Company


3.2 Odd-Chain

Vitamin B₁₂ Homolytic Cleavage




Catalyzes group migration


L-Methylmalonyl CoA



5'-Deoxyadenosyl radical



5'-Deoxyadenosine



Succinyl CoA




Figure 22-16
Biochemistry, Sixth Edition
© 2007 W.H. Freeman and Company

3.5 Ketone Bodies


Use of fatty acids in the citric acid cycle requires carbohydrates for the the production of oxaloacetate.

During starvation or diabetes, OAA is used to make glucose

- Fatty acids are then used to make ketone bodies (acetoacetate and D-3-hydroxybutarate)

3.5 Ketone Bodies

Ketone bodies, acetoacetate and 3-hydroxybutarate are formed from Acetyl-CoA



3.6 Ketone Bodies as a Fuel Source

The liver is the major source of ketone bodies.

- It is transported in the blood to other tissues

Acetoacetate in the tissues

- Acetoacetate is first activated to acetoacetyl-CoA by transferring the CoASH from succinyl-CoA.
- It is then split into two Acetyl-CoA by a thiolase reaction

3.6 Ketone Bodies as a Fuel Source




Figure 22-21
Biochemistry, Sixth Edition
 © 2007 W.H. Freeman and Company

3.6 Ketone Bodies as a Fuel Source





Figure 22-23
Biochemistry, Sixth Edition
© 2007 W. H. Freeman and Company

3.6 Ketone Bodies as a Fuel Source

Atkins Diet:

<http://www.webmd.com/diet/atkins-diet-what-it-is>


The measure of success?



3.7 Fatty Acids Cannot be Used to Synthesize Glucose


Even though the citric acid cycle intermediate oxaloacetate can be used to synthesize glucose, Acetyl-CoA cannot be used to synthesize oxaloacetate.

- The two carbons that enter the citric acid cycle as Acetyl-CoA leave as CO_2 .



Class problem

1. Explain the meaning (from a biochemistry perspective) of the saying “fats burn in the flame of carbohydrates.”



Update Alzheimer's Treatment

BBC
NEWS

[Watch](#) ONE-MINUTE WORLD NEWS



News Front Page



Africa

Americas

Asia-Pacific

Europe

Middle East

South Asia

UK

Business

Health

Medical notes

Science/Nature

Technology

Entertainment

Also in the news

Video and Audio

Have Your Say

In Pictures

Country Profiles

Special Reports

Page last updated at 00:20 GMT, Tuesday, 1 April 2008 01:20 UK

[E-mail this to a friend](#)

[Printable version](#)

Medication 'worsens Alzheimer's'

Anti-psychotic drugs commonly given to Alzheimer's patients often make their condition worse, a UK study suggests.

Neuroleptics provided no benefit for patients with mild behavioural problems, but were associated with a marked deterioration in verbal skills.

The research focused on 165 people with advanced Alzheimer's who were living in nursing homes in four British cities.

Up to 60% of Alzheimer's patients in nursing homes are given the drugs to control behaviour such as aggression.

The study appears in the journal Public Library of Science Medicine.

The researchers, from Kings College London and the Universities of Oxford and Newcastle, found the drugs offered no long-term benefit for most



CASE STUDY

Rita Clark's husband was diagnosed with Alzheimer's seven years ago. Rita, from Cleveland, said: "My

VIDEO AND AUDIO NEWS

What the report concluded

[Listen](#)

SEE ALSO

- ▶ 'Chemical cosh' early death risk
29 Mar 07 | Health
- ▶ Concerns over Alzheimer's drug
18 Feb 05 | Health
- ▶ Alzheimer's drug's impact hailed
14 May 07 | Health
- ▶ Alzheimer's disease
08 Jan 04 | Medical notes

RELATED INTERNET LINKS

- ▶ Alzheimer's Research Trust
- ▶ Public Library of Science Medicine
- ▶ Alzheimer's Society



4. Fatty Acid Synthesis.

Fatty acid are synthesized and degraded by different pathways.

- Synthesis takes place in the cytosol.
- Intermediates are attached to the acyl carrier protein (ACP).
- In higher organisms, the active sites for the synthesis reactions are all on the same polypeptide.
- The activated donor in the synthesis is malonyl-ACP.
- Fatty acid reduction uses $\text{NADPH} + \text{H}^+$.
- Elongation stops at C_{16} (palmitic acid)


4.1 Formation of Malonyl Coenzyme A

Formation of malonyl-CoA is the committed step in fatty acid synthesis.



4.2 Acyl Carrier Protein


The intermediates in fatty acid synthesis are covalently linked to the acyl carrier protein (ACP)



4.3 Elongation


In bacteria the enzymes that are involved in elongation are separate proteins; in higher organisms the activities all reside on the same polypeptide.

- To start an elongation cycle, Acetyl-CoA and Malonyl-CoA are each transferred to an acyl carrier protein



4.3 Elongation






Acyl-malonyl ACP
condensing enzyme forms
Acetoacetyl-ACP.



4.3 Elongation

The next three reactions are similar to the reverse of fatty acid degradation, except


- The NADPH is used instead of NADH and FADH_2
- The D-enantiomer of Hydroxybutyrate is formed instead of the L-enantiomer



4.3 Elongation

The elongation cycle is repeated six more times, using malonyl-CoA each time, to produce palmitoyl-ACP.

A thioesterase then cleaves the palmitoyl-CoA from the ACP.



4.4 Multifunctional Fatty Acid Synthase

Domain 1


- Substrate entry (AT & MT) and condensation unit (CE)

Domain 2


- Reduction unit (DH, ER & KR)


Domain 3

- Palmitate release unit (TE)




4.4 Multifunctional Fatty Acid Synthase





Distinct organization of two eukaryotic fatty acid synthases. (Left) Fungal fatty acid synthase assumes a barrel-like structure with multiple reaction chambers. (Right) Mammalian fatty acid synthase is a more compact, multi-domain complex with distinct active sites and reaction chambers. The acyl carrier protein of the mammalian enzyme carries malonyl, acetyl, and releases the palmitic acid. The acyl carrier protein of the mammalian enzyme is post-translationally modified by phosphopantetheinyl transferases that are likely localized as timers at the barrel apices.


4.5 Fatty Acid Synthase Mechanism




4.6 Stoichiometry of FA synthesis

The stoichiometry of palmitate synthesis:


- Synthesis of palmitate from Malonyl-CoA



- Synthesis of Malonyl-CoA from Acetyl-CoA



- Overall synthesis



4.7 Citrate Shuttle

Acetyl-CoA is synthesized in the mitochondrial matrix, whereas fatty acids are synthesized in the cytosol

- Acetyl-CoA units are shuttled out of the mitochondrial matrix as citrate:








Figure 22-28
 Biochemistry, Sixth Edition
 © 2007 W.H. Freeman and Company



4.8 Sources of NADPH

The malate dehydrogenase and NADP⁺-linked malate enzyme reactions of the citrate shuttle exchange NADH for NADPH



5. Regulation of Fatty Acid Synthesis


Regulation of Acetyl carboxylase

- Global



- + insulin
- - glucagon
- - epinephrine

- Local

- + Citrate
- - Palmitoyl-CoA
- - AMP




5.1 Regulation of Fatty Acid Synthesis



6. Elongation and Unsaturation

Endoplasmic reticulum systems introduce double bonds into long chain acyl-CoA's


- Reaction combines both NADH and the acyl-CoA's to reduce O_2 to H_2O .



6.1 Elongation and Unsaturation

Elongation and unsaturation convert palmitoyl-CoA to other fatty acids.


- Reactions occur on the cytosolic face of the endoplasmic reticulum.
- Malonyl-CoA is the donor in elongation reactions




6.2 Eicosanoid Hormones

Eicosanoid hormones are synthesized from arachadonic acid (20:4).


- Prostaglandins
 - 20-carbon fatty acid containing 5-carbon ring
 - Prostacyclins
 - Thromboxanes
- Leukotrienes
 - contain three conjugated double bonds




6.2 Eicosanoid Hormones




6.2 Eicosanoid Hormones




6.2 Eicosanoid Hormones




Prostaglandin A₂



Prostacyclin (PGI₂)



Thromboxane A₂ (TXA₂)



Leukotriene B₄

Class problem

1. Explain the meaning (from a biochemistry perspective) of the saying “fats burn in the flame of carbohydrates.” How would proteins fit into this statement?

