


Chem 150, Spring 2015

Unit 12 - Lipids

Introduction


- Biomolecules include
 - ✦ Proteins
 - ✦ Nucleic acids
 - ✦ Carbohydrates
 - ✦ **Lipids**
- Lipids are not defined by the chemical structures, like the other classes of biomolecules, but instead by their physical properties.
 - ✦ They are the components of cells that dissolve in non-polar (oily) solvents, and include fats, steroids, isoprenoids, phospholipids, glycolipids...



Chem 150, Unit 12: Lipids 2

Introduction


- Biomolecules include
 - ✦ Proteins
 - ✦ Nucleic acids
 - ✦ Carbohydrates
 - ✦ **Lipids**
- Lipids are not defined by the chemical structures, like the other classes of biomolecules, but instead by their physical properties.
 - ✦ They are the components of cells that dissolve in non-polar (oily) solvents, and include fats, steroids, isoprenoids, phospholipids, glycolipids...




Chem 150, Unit 12: Lipids 2

16.1 Fatty Acids and Triglycerides

- Animal fats and vegetable oils are triglycerides (triacylglycerols).
- Triglycerides are an important energy source tissues such as muscle, liver and kidneys.
- Triglycerides are made from one glycerol molecule plus three fatty acids (long chain carboxylic acid).




Glycerol
(glycerin)



Chem 150, Unit 12: Lipids 3

Triglycerides


- Triglycerides are formed by three condensation reactions.



Chem 150, Unit 12: Lipids 4

Saturated and Unsaturated Fatty Acids


- Saturated fatty acids** do not contain any alkene groups.
- Unsaturated fatty acids** contain at least one alkene group. Naturally occurring fatty acids contain *cis* alkene(s).



Chem 150, Unit 12: Lipids 5

Classification


- Monounsaturated fatty acid: contain only one alkene group
- Polyunsaturated fatty acid: contains more than one alkene group.
- Unsaturated fatty acids are classified by the number and position of first alkene group from the final carbon of the chain (omega carbon).



Chem 150, Unit 12: Lipids 6

Classification


- Monounsaturated fatty acid: contain only one alkene group
- Polyunsaturated fatty acid: contains more than one alkene group.
- Unsaturated fatty acids are classified by the number and position of first alkene group from the final carbon of the chain (omega carbon).



Chem 150, Unit 12: Lipids 6

Classification

- Monounsaturated fatty acid: contain only one alkene group
- Polyunsaturated fatty acid: contains more than one alkene group.
- Unsaturated fatty acids are classified by the number and position of first alkene group from the final carbon of the chain (omega carbon).



Chem 150, Unit 12: Lipids 6

Fatty Acids in Triglycerides

- Triglycerides often contain three different fatty acids, with any combination of saturated or unsaturated.
- Because fatty acids are a major component in triglycerides, fatty acids and triglycerides have similar physical properties. For example, they are both insoluble in water but soluble in organic solvents.

Chem 150, Unit 12: Lipids 7

Physical Properties


- Saturated fats are able to align themselves and tend to form solids at room temperature, therefore they have higher melting points.
- The cis alkene bonds of unsaturated fats create “kinks” in the structure that prevent close packing.
 - These “kinks” lower the melting points of fatty acids and the fats or oils that contain them.

Melting point =

Chem 150, Unit 12: Lipids 8

Physical Properties

- Saturated fats are able to align themselves and tend to form solids at room temperature, therefore they have higher melting points.
- The cis alkene bonds of unsaturated fats create “kinks” in the structure that prevent close packing.
 - These “kinks” lower the melting points of fatty acids and the fats or oils that contain them.



Melting point =

Chem 150, Unit 12: Lipids 8

Physical Properties

- Saturated fats are able to align themselves and tend to form solids at room temperature, therefore they have higher melting points.
- The cis alkene bonds of unsaturated fats create “kinks” in the structure that prevent close packing.
 - ✦ These “kinks” lower the melting points of fatty acids and the fats or oils that contain them.

Melting point =

Chem 150, Unit 12: Lipids 8

Physical Properties

- Saturated fats are able to align themselves and tend to form solids at room temperature, therefore they have higher melting points.
- The cis alkene bonds of unsaturated fats create “kinks” in the structure that prevent close packing.
 - ✦ These “kinks” lower the melting points of fatty acids and the fats or oils that contain them.

Melting point = 69°C

Chem 150, Unit 12: Lipids 8

Physical Properties

- Saturated fats are able to align themselves and tend to form solids at room temperature, therefore they have higher melting points.
- The cis alkene bonds of unsaturated fats create “kinks” in the structure that prevent close packing.
 - ✦ These “kinks” lower the melting points of fatty acids and the fats or oils that contain them.

Melting point = 69°C 14°C

Chem 150, Unit 12: Lipids 8

Physical Properties

- Saturated fats are able to align themselves and tend to form solids at room temperature, therefore they have higher melting points.
- The cis alkene bonds of unsaturated fats create “kinks” in the structure that prevent close packing.
 - ✦ These “kinks” lower the melting points of fatty acids and the fats or oils that contain them.

Melting point = 69°C 14°C -5°C

Chem 150, Unit 12: Lipids 8

Melting Points

- Percentage of saturated and unsaturated fats in common fats and oils effect the melting points.

TABLE 16.2 The Physical Properties and Fatty Acid Composition of Some Fats and Oils

Fat or Oil	Melting Point	State at Room Temperature	Saturated Fatty Acids	Monounsaturated Fatty Acids	Polyunsaturated Fatty Acids
Lard (pork fat)	34°C	Solid	41%	47%	12%
Butterfat	32°C	Solid	68%	28%	4%
Coconut oil	25°C	Solid	92%	8%	0%
Cod liver oil	-5°C	Liquid	20%	62%	18%
Olive oil	-6°C	Liquid	15%	75%	10%
Corn oil	-20°C	Liquid	13%	29%	58%

Melting Points


- Humans cannot synthesize polyunsaturated fatty acids, so these are **essential fatty acids** in our diet.

TABLE 16.1 The Structures of Common Fatty Acids

Name and Melting Point	Structure
Lauric acid (44°C)	$\text{CH}_3 - (\text{CH}_2)_{10} - \text{COOH}$
Myristic acid (55°C)	$\text{CH}_3 - (\text{CH}_2)_{12} - \text{COOH}$
Palmitic acid (63°C)	$\text{CH}_3 - (\text{CH}_2)_{14} - \text{COOH}$
Stearic acid (69°C)	$\text{CH}_3 - (\text{CH}_2)_{16} - \text{COOH}$
Palmitoleic acid (0°C)	$\text{CH}_3 - (\text{CH}_2)_5 - \text{CH} = \text{CH} - (\text{CH}_2)_7 - \text{COOH}$
Oleic acid (14°C)	$\text{CH}_3 - (\text{CH}_2)_7 - \text{CH} = \text{CH} - (\text{CH}_2)_7 - \text{COOH}$
Linoleic acid (-5°C)	$\text{CH}_3 - (\text{CH}_2)_4 - \text{CH} = \text{CH} - \text{CH}_2 - \text{CH} = \text{CH} - (\text{CH}_2)_7 - \text{COOH}$
Linolenic acid (-11°C)	$\text{CH}_3 - \text{CH}_2 - \text{CH} = \text{CH} - \text{CH}_2 - \text{CH} = \text{CH} - \text{CH}_2 - \text{CH} = \text{CH} - (\text{CH}_2)_7 - \text{COOH}$

16.2 Chemical Reactions of Triglycerides


- Unsaturated fatty acids can be hydrogenated to produce saturated fatty acids.



- A firm solid is produced when unsaturated triglycerides are hydrogenated.
- Saturated fats are more desirable in baking because they are more stable in the presence of oxygen and do not become rancid as rapidly.


16.2 Chemical Reactions of Triglycerides

- Often oils are only partially hydrogenated, and some of the *cis* fats are converted into *trans* fats in the process.




16.2 Chemical Reactions of Triglycerides

- Often oils are only partially hydrogenated, and some of the *cis* fats are converted into *trans* fats in the process.




16.2 Chemical Reactions of Triglycerides

- Often oils are only partially hydrogenated, and some of the *cis* fats are converted into *trans* fats in the process.




Hydrolysis of Triglycerides

- Triglycerides are esters and can be hydrolyzed.
- Hydrolysis by aqueous H_2SO_4 :




Hydrolysis by aqueous NaOH :




Fat Digestion

- Bile salts, which are like detergents, and enzymes are necessary to digest triglycerides.
- Bile salts break up (emulsify) clusters of fat molecules.
- Lipases**, which are enzymes, break the triglycerides into a monoglyceride (glycerol with a single fatty acid attached) and the conjugate bases of two fatty acids.



Digestion of a Triglyceride



Chem 150, Unit 12: Lipids 15

16.3 Catabolism of fatty acids

- Triglycerides are the most efficient energy sources: burning one gram of fat gives 9 kcal, carbohydrates or proteins give 4 kcal.
- Breakdown of fatty acids starts with an activation reaction (fatty acid \rightarrow fatty acyl-CoA) and then the four steps of beta oxidation.
- The addition of Coenzyme A allows for enzymes to recognize and bind to the fatty acid

Chem 150, Unit 12: Lipids 16


Beta Oxidation- Obtaining Energy from Fatty Acids

- Beta oxidation shortens the fatty acyl-CoA by two carbons, producing acetyl-CoA. The shortened fatty acyl-CoA then undergoes beta oxidation until all of the carbons have been used to produce acetyl-CoA in a series of 4 reactions

Chem 150, Unit 12: Lipids 17

Beta Oxidation- Obtaining Energy from Fatty Acids


- Beta oxidation shortens the fatty acyl-CoA by two carbons, producing acetyl-CoA. The shortened fatty acyl-CoA then undergoes beta oxidation until all of the carbons have been used to produce acetyl-CoA in a series of 4 reactions



Chem 150, Unit 12: Lipids 17

Beta Oxidation- Obtaining Energy from Fatty Acids

- Beta oxidation shortens the fatty acyl-CoA by two carbons, producing acetyl-CoA. The shortened fatty acyl-CoA then undergoes beta oxidation until all of the carbons have been used to produce acetyl-CoA in a series of 4 reactions



Chem 150, Unit 12: Lipids 17


Beta Oxidation- Obtaining Energy from Fatty Acids

- Beta oxidation shortens the fatty acyl-CoA by two carbons, producing acetyl-CoA. The shortened fatty acyl-CoA then undergoes beta oxidation until all of the carbons have been used to produce acetyl-CoA in a series of 4 reactions

Chem 150, Unit 12: Lipids 17

Reaction 1


- Dehydrogenation of the two carbons closest to the carbonyl group



Chem 150, Unit 12: Lipids 18

Reaction 2


- Hydration of the β -carbon



Chem 150, Unit 12: Lipids 19


Reaction 3

- Oxidation of the alcohol group




Reaction 4

- Carbon-carbon bond to the right of the ketone group is broken to produce acetyl-CoA and a new fatty acyl-CoA




Catabolism of Glucose



Three key features of beta oxidation

- It produces 1 each of NADH and FADH₂
- It forms a molecule of acetyl-CoA what can enter the citric acid cycle to form other high energy molecules
- It forms a new fatty acyl-CoA that can be used in another beta oxidation.

Beta Oxidation



Chem 150, Unit 12: Lipids 24

Example: Lauric Acid Oxidation

TABLE 16.3 The Complete Oxidation of Lauric Acid, a Saturated Fatty Acid

Pathway	Function	High-Energy Molecules Formed or Consumed*	Number of Occurrences When Lauric Acid Is Oxidized	Total High-Energy Molecules Formed or Consumed*
Activation	Converts lauric acid into lauryl-CoA	2 ATP broken down	1	2 ATP broken down
Beta oxidation	Breaks down lauryl-CoA into 6 molecules of acetyl-CoA	1 NADH 1 FADH ₂	5	5 NADH 5 FADH ₂
Citric acid cycle	Oxidizes the acetyl group of acetyl-CoA into CO ₂	1 ATP 3 NADH 1 FADH ₂	6	6 ATP 18 NADH 6 FADH ₂
Overall oxidation of lauric acid	Oxidizes lauric acid into CO ₂			2 ATP broken down 6 ATP formed 23 NADH 11 FADH ₂

*Consumed molecules are in red.

We have seen that the NADH + H⁺ and FADH₂ can go on to produce additional ATP's by coupling the electron transport chain, where these reduced nucleotides are deoxidized with O₂, with ATP synthase.

NADH + H⁺ will produce 2.5 ATP's and FADH₂ will produce 1.5 ATP's

Chem 150, Unit 12: Lipids 25

Example: Lauric Acid Oxidation

TABLE 16.3 The Complete Oxidation of Lauric Acid, a Saturated Fatty Acid

Pathway	Function	High-Energy Molecules Formed or Consumed*	Number of Occurrences When Lauric Acid Is Oxidized	Total High-Energy Molecules Formed or Consumed*
Activation	Converts lauric acid into lauryl-CoA	2 ATP broken down	1	2 ATP broken down
Beta oxidation	Breaks down lauryl-CoA into 6 molecules of acetyl-CoA	1 NADH 1 FADH ₂	5	5 NADH 5 FADH ₂
Citric acid cycle	Oxidizes the acetyl group of acetyl-CoA into CO ₂	1 ATP 3 NADH 1 FADH ₂	6	6 ATP 18 NADH 6 FADH ₂
Overall oxidation of lauric acid	Oxidizes lauric acid into CO ₂			2 ATP broken down 6 ATP formed 23 NADH 11 FADH ₂

*Consumed molecules are in red.

We have seen that the NADH + H⁺ and FADH₂ can go on to produce additional ATP's by coupling the electron transport chain, where these reduced nucleotides are deoxidized with O₂, with ATP synthase.

NADH + H⁺ will produce 2.5 ATP's and FADH₂ will produce 1.5 ATP's

Chem 150, Unit 12: Lipids 25

Unsaturated Fatty Acids

- Unsaturated fatty acids undergo a similar process, but produce one FADH₂ fewer than a saturated fatty acid for every double bond in the chain.

Chem 150, Unit 12: Lipids 26

Try It!

Question:

How many cycles of beta oxidation are required to break down one molecule of palmitic acid, a saturated fatty acid that has the molecular formula $C_{16}H_{32}O_2$? How many molecules of acetyl-CoA will be formed? What will the ATP yield be?

Fatty Acids produce similar amounts of ATP on a Mass Basis

- 1 mole Lauric acid (200.3 g/mol) produces 78 molecules of ATP and 1 mole palmitic acid (256.4 g/mol) produces 106 molecules of ATP.
- When comparing the breakdown of 100 g each by mass, you see they make very similar amounts of ATP:

$$100 \text{ g lauric acid} \left(\frac{78 \text{ mol ATP}}{1 \text{ mol lauric acid}} \right) \left(\frac{1 \text{ mol lauric acid}}{200.3 \text{ g lauric acid}} \right) = 38.9 \text{ mol ATP}$$
$$100 \text{ g palmitic acid} \left(\frac{106 \text{ mol ATP}}{1 \text{ mol palmitic acid}} \right) \left(\frac{1 \text{ mol palmitic acid}}{256.4 \text{ g palmitic acid}} \right) = 41.3 \text{ mol ATP}$$

Fats vs Carbohydrates

- Comparing this to carbohydrates

$$100 \text{ g glucose} \left(\frac{32 \text{ mol ATP}}{1 \text{ mol glucose}} \right) \left(\frac{1 \text{ mol glucose}}{180.2 \text{ g glucose}} \right) = 17.8 \text{ mol ATP}$$


$$\approx \frac{40 \text{ mol ATP from fat}}{18 \text{ mol ATP from carbohydrates}} = 2.2 \text{ times more energy from fat}$$

16.4 Glycerophospholipids and Cell Membranes

- Cell membranes are made of phospholipids, which contain one or more long hydrocarbon chain attached to a phosphate group.
- Example: Glycerophospholipids are composed of two fatty acids, glycerol, a phosphate group, and a polar molecule (typically choline, serine or ethanolamine).




General structure of a triglyceride



General structure of a glycerophospholipid


Formation of a Glycerophospholipid



Chem 150, Unit 12: Lipids 31

Lipid Bilayer


- Earlier we discussed how soaps aggregate to form micelles when mixed with water.
- Glycerophospholipids aggregate to form lipid bilayers when mixed with water



Chem 150, Unit 12: Lipids 32

Lipid Bilayer


- A lipid bilayer is the basic component of cell membranes.
- Hydrophilic groups of the glycerophospholipids face out to the aqueous intra- and extracellular environment. Hydrophobic tails face each other.
- This creates a strong flexible coating for a cell that prevents free exchange of molecules and ions between the inside and outside of a cell.



Chem 150, Unit 12: Lipids 33

Other Components in the Lipid Bilayer


- Membranes contain many other molecules, including proteins and steroids.




Chem 150, Unit 12: Lipids 34

Steroids and Cholesterol

- Steroids are a class of lipids that contain a fused ring nucleus.



- Cholesterol is a common steroid.




Chem 150, Unit 12: Lipids 35

Proteins in the Cell Membrane

- Many of the proteins in membranes are transport proteins
 - These proteins provide passageways for ions and molecules to cross the lipid bilayer.
- Passive transport occurs when a molecule or ion diffuses from higher concentration to lower concentration (down a **concentration gradient**).
- Active transport occurs when a molecule or ion moves from lower concentration to higher concentration (up a **concentration gradient**).
 - Active transport requires a source of energy.

Chem 150, Unit 12: Lipids 36

Active and Passive Transport



Chem 150, Unit 12: Lipids 37

16.5 Concentration gradients and ATP formation

- Prokaryotes (bacteria) have a less complex internal structure than eukaryotes.
 - Eukaryotic cells contain membrane bound organelles.


TABLE 16.7 Organelles in Eukaryotic Cells

Structure	Function
Endoplasmic reticulum	Builds proteins and lipids
Golgi complex	Processes and sorts proteins after they are synthesized
Lysosome	Hydrolyzes worn-out and damaged cell components
Peroxisome	Oxidizes waste products and foreign material
Mitochondrion	Carries out most catabolic pathways, and produces ATP for the cell
Chloroplast (plants only)	Carries out photosynthesis

Chem 150, Unit 12: Lipids 38

Mitochondria


- Eukaryotic cell.



Chem 150, Unit 12: Lipids 39

Mitochondria


- One of these, the mitochondria, is the primary location for ATP production in eukaryotic cells .



Chem 150, Unit 12: Lipids 40

Mitochondria

- One of these, the mitochondria, is the primary location for ATP production in eukaryotic cells .




Mitochondria are believed to have evolved from a free-living bacterium

Chem 150, Unit 12: Lipids 40

Electron Transport Chain

- A set of enzymes in the mitochondria carry out the electron transport chain. These enzymes use O_2 to convert $FADH_2$ and $NADH$ back to their oxidized forms and release energy:




- The energy is used to produce a concentration gradient of H^+ ion across the mitochondrial membrane, which is then used to produce ATP.

Chem 150, Unit 12: Lipids 41

Electron Transport Chain


- Energy is used by the electron transport chain to actively transport H^+ into the intermembrane space, creating a concentration gradient.



Chem 150, Unit 12: Lipids 42

Charge and Concentration Gradient

- In addition to a concentration gradient, a charge gradient is also produced because there is now a high concentration of positive ions in the intermembrane space.



Chem 150, Unit 12: Lipids 43


FADH₂ vs NADH

- More energy is released reoxidizing $NADH + H^+$ than reoxidizing $FADH_2$
 - Therefore, fewer H^+ are transported across the membrane when oxidizing $FADH_2$.
- The electron transport chain is able to push 10 H^+ through the membrane for every $NADH + H^+$ oxidized, but only 6 H^+ for every $FADH_2$ oxidized.

Chem 150, Unit 12: Lipids 44


ATP Synthase

- Next, hydrogen ions move through ATP synthase back into the matrix.
- This movement releases the energy from the gradient and allows ATP synthase to produce ATP from ADP and P_i . Movement of 3 H^+ are required to make one molecule ATP.




Chem 150, Unit 12: Lipids 45

Electron Transport Chain and ATP Synthase



Chem 150, Unit 11: Carbohydrates 46

Electron Transport Chain and ATP Synthase



Chem 150, Unit 11: Carbohydrates 46

Electron Transport Chain and ATP Synthase



Chem 150, Unit 11: Carbohydrates 46


Transport Proteins

- A transport protein then moves ATP into the intermembrane and one H^+ , ADP, P_i into the matrix.
- Movement of 4 H^+ **total** are required to make one molecule ATP
- In terms of coenzymes:
 - One mole of NADH produces 2.5 moles ATP
 - One mole of $FADH_2$ produces 1.5 moles ATP

Chem 150, Unit 12: Lipids 47

Oxidative Phosphorylation

- The process that begins with oxidation of NADH and FADH_2 and ends with ATP production and transport is known as oxidative phosphorylation.



Chem 150, Unit 12: Lipids 48

Other Uses for Concentration Gradients

- Concentration gradients supply energy for other purposes.
- Cells use ATP hydrolysis to create gradients with sodium and potassium ions, which in turn can be used for other cellular processes like nerve impulse transmission

Chem 150, Unit 12: Lipids 49

Other Uses for Concentration Gradients

- Unit 13: Nucleic Acids and Protein Synthesis
 - Chapter 17 in Armstrong

Chem 150, Unit 12: Lipids 50