

Chem 150, Spring 2015

Unit 6 - Alcohols, and Hydration & Dehydration Reactions

Introduction

Question:

What do the words hydration and dehydration mean to you?

Chem 150, Unit 6: Alcohols 2

10.1 The Hydration Reaction

- In the last unit we introduced the concept of functional groups.
- A functional group is a group of atoms the combine to give a molecule a characteristic set of chemical and physical properties.*

Chem 150, Unit 6: Alcohols 3

10.1 The Hydration Reaction

- In the last unit we introduced the concept of functional groups.

A functional group is a group of atoms the combine to give a molecule a characteristic set of chemical and physical properties.

TABLE 9.10 Some Representative Functional Groups in Organic Chemistry

Functional Group	Name	An Example of a Compound That Contains This Group:
$\text{C}=\text{C}$	alkene	$\text{CH}_3-\text{CH}=\text{CH}_2$ propene
$\text{C}\equiv\text{C}$	alkyne	$\text{CH}_3-\text{C}\equiv\text{CH}$ propyne
$\text{C}-\text{Cl}$	chloroalkane	$\text{CH}_3-\text{CH}_2-\text{CH}_2-\text{Cl}$ 1-chloropropane
$\text{C}-\text{O}-\text{H}$	alcohol	$\text{CH}_3-\text{CH}_2-\text{CH}_2-\text{OH}$ 1-propanol
$\text{C}-\text{N}-\text{H}$	amine	$\text{CH}_3-\text{CH}_2-\text{CH}_2-\text{NH}_2$ propylamine
$\text{O}=\text{C}-\text{H}$	aldehyde	$\text{CH}_3-\text{CH}_2-\text{C}(=\text{O})-\text{H}$ propanal
$\text{O}=\text{C}-\text{O}-\text{H}$	carboxylic acid	$\text{CH}_3-\text{CH}_2-\text{C}(=\text{O})-\text{OH}$ propanoic acid

Chem 150, Unit 6: Alcohols 3

10.1 The Hydration Reaction


- In the last unit we introduced the concept of functional groups.
 - A functional group is a group of atoms the combine to give a molecule a characteristic set of chemical and physical properties.

10.1 The Hydration Reaction


- In the last unit we introduced the concept of functional groups.
 - A functional group is a group of atoms the combine to give a molecule a characteristic set of chemical and physical properties.
 - For example, the **alkene** functional group can react with water to produce the **alcohol** functional group

10.1 The Hydration Reaction

- An reaction that alkenes undergo in biological chemistry is the hydration reaction.
- The product of a hydration reaction is called an alcohol and contains a hydroxyl group.




Many Alkenes Produce Two Hydration Products




Many Alkenes Produce Two Hydration Products

- Alkenes can also participate in other addition type reactions




Markovnikov's Rule

- If there is an option, *the hydrogen atom of water prefers to become attached to the alkene carbon that is bonded to the greater number of hydrogen atoms.*




Cycloalkenes Can Be Hydrated

- Cycloalkenes can also react with water to form alcohols.





Cycloalkenes Can Be Hydrated

- Cycloalkenes can also react with water to form alcohols.




Cycloalkenes Can Be Hydrated

- Cycloalkenes can also react with water to form alcohols.



Chem 150, Unit 6: Alcohols 8

Symmetrical Alkenes Produce Only One Hydration Product





Chem 150, Unit 6: Alcohols 9

Try It!

Question:

Draw the structure(s) of the hydration products for each of the following alkenes.




Chem 150, Unit 6: Alcohols 10

10.2 Controlling the Product: An Introduction to Enzymes

- Living organisms carry out many reactions. These reactions occur in specific sequences, in which each product becomes the starting material for the next reaction.
- Living organisms solve the problem of multiple products using enzymes.
 - Enzymes are biological catalysts that produce a single product.
 - Enzymes are also selective about the reactants that they can use so they will only react with certain compounds.


Chem 150, Unit 6: Alcohols 11

Enzyme Selectivity in a Hydration Reaction



Chem 150, Unit 6: Alcohols 12

Enzyme Selectivity in a Hydration Reaction



Chem 150, Unit 6: Alcohols 13

10.3 Naming Alcohols




1. Name the hydrocarbon framework.
2. Identify the functional group by modifying the ending of the alkane name. For alcohols, change the end of the name of the corresponding hydrocarbon from *-e* to *-ol*.
3. To complete the root name, add a number to tell where the functional group is located. Number from the end of the carbon chain closest to the hydroxyl group.
4. Identify and locate any alkyl groups and append their names to the front of the root name.

Chem 150, Unit 6: Alcohols 14

Try It!

Questions:

Give the IUPAC name for the following molecules



Chem 150, Unit 5: Hydrocarbons 15


Try It!

Questions:

Draw structures for the following named molecules.


- 3-isopropyl-1-nonol
- 2-propanol

Commonly Used Trivial names



10.4 The Physical Properties of Alcohols

- The two covalent bonds in the alcohol functional group are strongly polar.
- The polar O-H bond allows alcohols to form hydrogen bonds.
- The ability to form hydrogen bonds gives alcohols quite different physical properties from those of hydrocarbons.



Boiling Points of Some Alkanes and Alcohols

TABLE 10.1 The Physical Properties of Some Alkanes and Alcohols

Alkane	Boiling Point	State at Room Temperature	Alcohol	Boiling Point	State at Room Temperature
Methane	-161°C	Gas	Methanol	65°C	Liquid
Butane	-1°C	Gas	1-Butanol	117°C	Liquid
Octane	126°C	Liquid	1-Octanol	194°C	Liquid

Boiling Points of Some Alkanes and Alcohols

- The addition of a hydroxyl group (-OH) greatly increases the boiling point of a hydrocarbon.

TABLE 10.1 The Physical Properties of Some Alkanes and Alcohols

Alkane	Boiling Point	State at Room Temperature	Alcohol	Boiling Point	State at Room Temperature
Methane	-161°C	Gas	Methanol	65°C	Liquid
Butane	-1°C	Gas	1-Butanol	117°C	Liquid
Octane	126°C	Liquid	1-Octanol	194°C	Liquid

Boiling Points of Some Alkanes and Alcohols

- The addition of a hydroxyl group (-OH) greatly increases the boiling point of a hydrocarbon.

TABLE 10.1 The Physical Properties of Some Alkanes and Alcohols

Alkane	Boiling Point	State at Room Temperature	Alcohol	Boiling Point	State at Room Temperature
Methane	-161°C	Gas	Methanol	65°C	Liquid
Butane	-1°C	Gas	1-Butanol	117°C	Liquid
Octane	126°C	Liquid	1-Octanol	194°C	Liquid

Boiling Points of Some Alkanes and Alcohols

- The addition of a hydroxyl group (-OH) greatly increases the boiling point of a hydrocarbon.
- Note the affect that dispersion interactions still have on the boiling points of alcohols

TABLE 10.1 The Physical Properties of Some Alkanes and Alcohols

Alkane	Boiling Point	State at Room Temperature	Alcohol	Boiling Point	State at Room Temperature
Methane	-161°C	Gas	Methanol	65°C	Liquid
Butane	-1°C	Gas	1-Butanol	117°C	Liquid
Octane	126°C	Liquid	1-Octanol	194°C	Liquid

Boiling Points of Some Alkanes and Alcohols


- The addition of a hydroxyl group (-OH) greatly increases the boiling point of a hydrocarbon.
- Note the affect that dispersion interactions still have on the boiling points of alcohols

TABLE 10.1 The Physical Properties of Some Alkanes and Alcohols

Alkane	Boiling Point	State at Room Temperature	Alcohol	Boiling Point	State at Room Temperature
Methane	-161°C	Gas	Methanol	65°C	Liquid
Butane	-1°C	Gas	1-Butanol	117°C	Liquid
Octane	126°C	Liquid	1-Octanol	194°C	Liquid


Hydrophobic and Hydrophilic Regions

- The solubility of a molecular compound depends on the relative sizes of the hydrophilic (-OH) and hydrophobic regions (CH₃CH₂-...) in the molecule.



Hydrophobic region

Methanol has a small hydrophobic region, so it has a high solubility in water.



Hydrophobic region

1-Hexanol has a large hydrophobic region, so it has a low solubility in water.

Hydrophobic and Hydrophilic Regions


- The solubility of a molecular compound depends on the relative sizes of the hydrophilic (-OH) and hydrophobic regions (CH₃CH₂-...) in the molecule.

TABLE 10.2 The Solubilities of Some Alcohols

Compound	Carbon Atoms	Solubility in 100 g of Water
Methanol	One	No limit
Ethanol	Two	(any amount of these alcohols will mix with water)
1-Propanol	Three	
1-Butanol	Four	7.4 g
1-Pentanol	Five	2.7 g
1-Hexanol	Six	0.7 g
1-Heptanol	Seven	0.1 g


Hydrophobic and Hydrophilic Regions

- The solubility of a molecular compound depends on the relative sizes of the hydrophilic (-OH) and hydrophobic regions (CH₃CH₂-...) in the molecule.



Hydrophobic region

Methanol has a small hydrophobic region, so it has a high solubility in water.



Hydrophobic region

1-Hexanol has a large hydrophobic region, so it has a low solubility in water.

Hydrophobic and Hydrophilic Regions


- The solubility of a molecular compound depends on the relative sizes of the hydrophilic (-OH) and hydrophobic regions (CH₃CH₂-...) in the molecule.

TABLE 10.3 The Dependence of Solubility on the Number of Hydroxyl Groups

Compound	OH Groups	Solubility in 100 g of H ₂ O
CH ₃ -CH ₂ -CH ₂ -CH ₂ -CH ₂ -CH ₃	None	0.04 g
OH CH ₂ -CH ₂ -CH ₂ -CH ₂ -CH ₂ -CH ₃	One	0.7 g
OH CH ₂ -CH ₂ -CH ₂ -CH ₂ -CH ₂ -CH ₂ -OH	Two	6 g
OH OH OH CH ₂ -CH ₂ -CH-CH ₂ -CH ₂ -CH ₂	Three	No limit


Hydrophobic and Hydrophilic Regions

- The solubility of a molecular compound depends on the relative sizes of the hydrophilic (-OH) and hydrophobic (CH₃CH₂-...) in the molecule.



Hydrophobic region

Methanol has a small hydrophobic region, so it has a high solubility in water.



Hydrophobic region

1-Hexanol has a large hydrophobic region, so it has a low solubility in water.

Solubility Trends


- Compounds with a hydrogen-bonding group (hydrophilic) dissolve better than compounds that cannot form hydrogen bonds, regardless of the sizes of the molecules.
- If two compounds have the same hydrophilic group, the molecule with the *smaller carbon framework* is the more soluble.
- If two compounds have the same carbon framework, the molecule with *more hydrophilic groups* is the more soluble.

Organic Compounds Generally Mix

- There is a wide range of common organic solvents, and there is no simple way to work out which compounds dissolve in which solvents.
- Most organic compounds dissolve reasonably well in organic solvents.
- Only compounds that approach water in their ability to form hydrogen bonds do not dissolve in hydrocarbons.
 - They do, however, dissolve well in alcohols because they can hydrogen to the solvent as well.



Alcohols Are Not Acidic or Basic

- Alcohols do not contain hydroxide ions and solutions in water are not appreciably acidic or basic.



Alcohols Are Not Acidic or Basic



- The hydroxyl group is covalently bonded to a carbon atom in an alcohol, and it does not dissociate from the rest of the molecule when the alcohol dissolves in water.
- † This is not true of carboxylic acid groups, which can dissociate in water.



Chem 150, Unit 6: Alcohols 24

Alcohols Are Not Acidic or Basic



- The hydroxyl group is covalently bonded to a carbon atom in an alcohol, and it does not dissociate from the rest of the molecule when the alcohol dissolves in water.
- † This is not true of carboxylic acid groups, which can dissociate in water.



Chem 150, Unit 6: Alcohols 24

Alcohols Are Not Acidic or Basic

- The hydroxyl group is covalently bonded to a carbon atom in an alcohol, and it does not dissociate from the rest of the molecule when the alcohol dissolves in water.
- † This is not true of carboxylic acid groups, which can dissociate in water.



Chem 150, Unit 6: Alcohols 24

10.5 Chirality in Organic Molecules

- A chiral object is an object that cannot be superimposed on its mirror image.
- † The same relationship can occur for some organic molecules.

Chem 150, Unit 6: Alcohols 25

10.5 Chirality in Organic Molecules

- A chiral object is an object that cannot be superimposed on its mirror image.
 - ✦ The same relationship can occur for some organic molecules.



A left hand is chiral, because it cannot be superimposed on its mirror image (a right hand).

Chem 150, Unit 6: Alcohols 25

10.5 Chirality in Organic Molecules

- A chiral object is an object that cannot be superimposed on its mirror image.
 - ✦ The same relationship can occur for some organic molecules.



A nose is achiral, because it is identical to its mirror image.

Chem 150, Unit 6: Alcohols 25

10.5 Chirality in Organic Molecules

- A chiral object is an object that cannot be superimposed on its mirror image.
 - ✦ The same relationship can occur for some organic molecules.



A left hand is chiral, because it cannot be superimposed on its mirror image (a right hand).





A nose is achiral, because it is identical to its mirror image.

Chem 150, Unit 6: Alcohols 25

Enantiomers

- The two mirror-image forms of a chiral molecule are called enantiomers.

One form of 2-butanol looks like this:




The other form of 2-butanol is the mirror image of the first form.

Chem 150, Unit 6: Alcohols 26


Chirality and Biomolecules

- Many important substances in medicine and biochemistry are chiral, including all proteins, most fats, all common carbohydrates, cholesterol, and a range of medications.
- All enzymes are chiral molecules, so they can distinguish between the two forms of other chiral molecules.
- Our bodies can normally use only one of the two forms of a chiral compound. The other form has little or no activity, and it may even be harmful.


Chirality and Biomolecules



Chirality and Biomolecules




Chirality and Biomolecules



Chirality and Biomolecules

- Receptor protein in our nose can distinguish between the two enantiomers of chiral molecule Carvone




Identifying Chiral Molecules

- Chiral molecules contain a chiral carbon. A carbon is chiral when it is bonded to four different groups of atoms.
- Many important molecules in biology and medicine contain more than one chiral carbon atom. All molecules that contain just one chiral carbon atom are chiral, but molecules that have two or more chiral carbon atoms are not necessarily chiral.
- You should be able to identify a chiral carbon.


Identifying Chiral Molecules

- Chiral molecules contain a chiral carbon. A carbon is chiral when it is bonded to four different groups of atoms.



Identifying Chiral Molecules


- Chiral molecules contain a chiral carbon. A carbon is chiral when it is bonded to four different groups of atoms.



- You should be able to identify a chiral carbon.

Identifying Chiral Molecules

- Chiral molecules contain a chiral carbon. A carbon is chiral when it is bonded to four different groups of atoms.



Identifying Chiral Molecules

- Chiral molecules contain a chiral carbon. A carbon is chiral when it is bonded to four different groups of atoms.
- Many important molecules in biology and medicine contain more than one chiral carbon atom. All molecules that contain just one chiral carbon atom are chiral, but molecules that have two or more chiral carbon atoms are not necessarily chiral.
- You should be able to identify a chiral carbon.

Try It!


Question

Draw the structures for the following biological compounds and circle the chiral carbons?

- A. lactic acid
- B. glucose
- C. glyceraldehyde


10.6 The Dehydration Reaction

- A dehydration reaction is one of the means by which our bodies remove oxygen from organic compounds.
- In the dehydration of an alcohol, the hydroxyl group and a hydrogen atom from an adjacent carbon are removed to make water and an alkene.




The Dehydration of 1-Propanol



When we dehydrate 1-propanol, we remove the OH group from carbon #1...




... so we must remove a hydrogen atom from the neighboring carbon (carbon #2).



These hydrogen atoms are not attached to the neighboring carbon, so they cannot be removed in this dehydration reaction.




Remove OH and H




These two carbon atoms need one more bond.

Add a bond between the two carbon atoms.




Many Dehydration Reactions Produce More Than One Product

- If an alcohol has more than one carbon atom next to the functional group, it can usually form more than one dehydration product.




Removing H from carbon #1 produces 1-butene.




Removing H from carbon #2 produces 2-butene (both the *cis* and *trans* isomers).

The Dehydration of Cyclic Alcohols


- Cyclic alcohols can also be dehydrated.




Remove H and OH




Product #1



Remove H and OH



Product #2



Line structures of the products

More on Dehydration Reactions


- Some alcohols do not have any hydrogen atoms attached to the adjacent carbon atoms and cannot be dehydrated.
- In our bodies, enzymes catalyze dehydration reactions.
 - Enzymes only make one alkene product when they dehydrate an alcohol.
 - This is important because the alkene products made are the specific ones your body needs for subsequent reactions.

More on Dehydration Reactions

ENZYME



The enzyme holds the water molecule in place so it can only react with the alkene group in the correct way.



Chem 150, Unit 6: Alcohols 37

More on Dehydration Reactions

ENZYME



The enzyme holds the water molecule in place so it can only react with the alkene group in the correct way.



Chem 150, Unit 6: Alcohols 37

10.7 Phenols and Thiols

- A phenol is a compound that contains a hydroxyl group bonded to a benzene ring.
 - Phenols cannot be dehydrated like alcohols.
 - Phenols cannot be formed by adding water to an alkene like alcohols.
 - Phenols are weak acids, unlike alcohols which are neither acidic or basic.
 - The hydroxyl group in phenols can hydrogen bond just like the hydroxyl group in alcohols.

Chem 150, Unit 6: Alcohols 38

The Properties of a Phenol, an Alcohol, and a Hydrocarbon

TABLE 10.5 The Properties of a Phenol, an Alcohol, and a Hydrocarbon

	Phenol	Cyclohexanol	Benzene
Structure			
Boiling point	182°C	161°C	80°C
Solubility in water	65 g/L	36 g/L	0.6 g/L
pH of 1% solution	5.5 (weakly acidic)	7.0 (neutral)	We cannot make a 1% solution.




Chem 150, Unit 6: Alcohols 39

The Properties of a Phenol, an Alcohol, and a Hydrocarbon

Question

Why are the boiling points for phenol and cyclohexanol substantially higher than that for benzene?

TABLE 10.5 The Properties of a Phenol, an Alcohol, and a Hydrocarbon

	Phenol	Cyclohexanol	Benzene
Structure			
Boiling point	182°C	161°C	80°C
Solubility in water	65 g/L	36 g/L	0.6 g/L
pH of 1% solution	5.5 (weakly acidic)	7.0 (neutral)	We cannot make a 1% solution.

Thiols

- Thiols are related to alcohols because a thiol is like a hydroxyl group where the oxygen is replaced with a sulfur.
 - ✦ The thiol group cannot form hydrogen bonds, so thiols have lower boiling points and evaporate more readily than alcohols and are less soluble in water.
 - ✦ The thiol group is weakly polar so it is attracted to water molecules. Thiols are more soluble in water than similar-sized alkanes.

The Properties of an Alcohol, a Thiol, and an Alkane

TABLE 10.6 The Properties of an Alcohol, a Thiol, and an Alkane

	Ethanol	Ethanethiol	Propane
Structure	$\text{CH}_3\text{—CH}_2\text{—OH}$	$\text{CH}_3\text{—CH}_2\text{—SH}$	$\text{CH}_3\text{—CH}_2\text{—CH}_3$
Boiling point	78°C	35°C	−42°C
Solubility in water	No limit	7 g/L	0.1 g/L

What Stinks?

- Thiols are known for their terrible odors.
- Low-molecular-weight thiols have some of the most offensive aromas in all of chemistry.
 - ✦ Examples: The odor of a skunk or freshly cut onions.
- Natural gas suppliers add a tiny amount of a thiol to natural gas (which has no odor) so leaks will be noticed and reported.
- An example of an exception (a good smell): a thiol is responsible for the pleasant aroma of grapefruit, coffee and garlic.

Next Up

- Unit 7: Carbonyls, Aldehydes and Ketons
 - ✦ Chapter 11 in Armstrong
 - ✦ Unit 7 Assignments due on 12. March, with deadline of 19. March.
