


Chem 352 - Lecture 7

Introduction to Metabolism

Question for the Day: How is metabolism like a flowing river?



1-1



Chem 352 - Lecture 7

Introduction to Metabolism

Question for the Day: How is metabolism like a flowing river?

Or as Heraclitus (c. 535 - 475 BCE) said

"Πάντα χωρεῖ καὶ οὐδὲν μένει" καὶ "οἷς ἐς τὸν αὐτὸν ποταμὸν οὐκ ἂν ἐμβαίῃς"

1-2

Chem 352 - Lecture 7



Introduction to Metabolism

Question for the Day: How is metabolism like a flowing river?

Or as Heraclitus (c. 535 - 475 BCE) said

"Πάντα χωρεῖ καὶ οὐδὲν μένει" καὶ "οἷς ἐς τὸν αὐτὸν ποταμὸν οὐκ ἂν ἐμβαίῃς"

"Everything changes and nothing remains still ... and ... you cannot step twice into the same stream"





1-3

Introduction

Metabolism is the sum total of all of the reactions that take place in a living cell.

- Made up of
 - metabolites, connected to each other by
 - enzyme catalyzed reactions




Chem 352, Lecture 7, Introduction to Metabolism 2

2-1

Introduction


Metabolism is the sum total of all of the reactions that take place in a living cell.

- Made up of
 - metabolites, connected to each other by
 - enzyme catalyzed reactions




Chem 352, Lecture 7, Introduction to Metabolism 2


2-2




2-3




2-4




2-5




3-1




3-2



3-3




4-1



4-2



5-1




5-2

Metabolic Pathways

Common themes in metabolism:

- Organisms or cells maintain specific internal




Chem 352, Lecture 7, Introduction to Metabolism 5

5-3

Metabolic Pathways

Common themes in metabolism:

- Organisms or cells maintain specific internal concentrations of inorganic ions, metabolites, and enzymes. Cell membranes provide the physical barrier that segregates cell components from the environment.




Chem 352, Lecture 7, Introduction to Metabolism 5

5-4

Metabolic Pathways

Common themes in metabolism:




Chem 352, Lecture 7, Introduction to Metabolism 6

6-1

Metabolic Pathways

Common themes in metabolism:

- Organisms extract energy from external sources to drive energy-consuming reactions. Photosynthetic organisms derive energy from the conversion of solar energy to chemical energy. Other organisms obtain energy from the ingestion and catabolism of energy-yielding compounds




Chem 352, Lecture 7, Introduction to Metabolism 6

6-2

Metabolic Pathways

Common themes in metabolism:




Metabolism 6

6-3

Metabolic Pathways

Common themes in metabolism:

- Organisms extract energy from external sources to drive energy-consuming reactions. Photosynthetic organisms derive energy from the conversion of solar energy to chemical energy. Other organisms obtain energy from the ingestion and catabolism of energy-yielding compounds



Chem 352, Lecture 7, Introduction to Metabolism 6

6-4

Metabolic Pathways

Common themes in metabolism:




Chem 352, Lecture 7, Introduction to Metabolism 7

7-1

Metabolic Pathways

Common themes in metabolism:

- The metabolic pathways in each organism are specified by the genes it contains in its genome.




Chem 352, Lecture 7, Introduction to Metabolism 7

7-2

Metabolic Pathways

Common themes in metabolism:




Chem 352, Lecture 7, Introduction to Metabolism 7

7-3

Metabolic Pathways

Common themes in metabolism:

- The metabolic pathways in each organism are specified by the genes it contains in its genome.



Chem 352, Lecture 7, Introduction to Metabolism 7

7-4

Metabolic Pathways

Common themes in metabolism:




Chem 352, Lecture 7, Introduction to Metabolism 8

8-1

Metabolic Pathways

Common themes in metabolism:

- Organisms and cells interact with their environment. The activities of cells must be geared to the availability of energy. Organisms grow and reproduce when the supply of energy from the environment is plentiful. When the supply of energy from the environment is limited, energy demands can be temporarily met by using internal stores or by slowing metabolic rates as in hibernation, sporulation, or seed formation. If the shortage is prolonged, organisms die.



Chem 352, Lecture 7, Introduction to Metabolism 8

8-2

Metabolic Pathways

Common themes in metabolism:




Chem 352, Lecture 7, Introduction to Metabolism 9

9-1

Metabolic Pathways

Common themes in metabolism:

- The cells of organisms are not static assemblies of molecules. Many cell components are continually synthesized and degraded, that is, they undergo turnover, even though their concentrations may remain virtually constant. The concentrations of other compounds change in response to changes in external or internal conditions.



Chem 352, Lecture 7, Introduction to Metabolism 9

9-2



9-3

Metabolic Pathways

Common themes in metabolism:


- The cells of organisms are not static assemblies of molecules. Many cell components are continually synthesized and degraded, that is, they undergo turnover, even though their concentrations may remain virtually constant. The concentrations of other compounds change in response to changes in external or internal conditions.

Chem 352, Lecture 7, Introduction to Metabolism 9

9-4

Metabolic Pathways

The enzymes arrange the metabolites into pathways.




Chem 352, Lecture 7, Introduction to Metabolism 10

10-1

Metabolic Pathways

The enzymes arrange the metabolites into pathways.




Chem 352, Lecture 7, Introduction to Metabolism 10

10-2

Metabolic Pathways

The enzymes arrange the metabolites into pathways.




Chem 352, Lecture 7, Introduction to Metabolism 10

10-3

Metabolic Pathways

The enzymes arrange the metabolites into pathways.



Chem 352, Lecture 7, Introduction to Metabolism 10

10-4

Metabolic Pathways

The enzymes arrange the metabolites into pathways.




Chem 352, Lecture 7, Introduction to Metabolism 10

10-5

Metabolic Pathways

The enzymes arrange the metabolites into pathways.



Chem 352, Lecture 7, Introduction to Metabolism 10

10-6

Metabolic Pathways

The enzymes arrange the metabolites into pathways.



Chem 352, Lecture 7, Introduction to Metabolism 10

10-7

Metabolic Pathways

Why pathways are broken down into little steps.

- Enzyme specificity allows only for breakage or formation of a few bonds at a time.
- Using pathways allows for multiple entry and exit points for metabolites.
- Pathways allow for finer control of energy input and output.
- Reactions are thermodynamically more efficient if carried out near equilibrium.


Chem 352, Lecture 7, Introduction to Metabolism 11

11-1

Metabolic Pathways

Why pathways are broken down into little steps.

- Enzyme specificity allows only for breakage or formation of a few bonds at a time.
- Using pathways allows for multiple entry and exit points for metabolites.
- Pathways allow for finer control of energy input and output.
- Reactions are thermodynamically more efficient if carried out near equilibrium.



Chem 352, Lecture 7, Introduction to Metabolism 11

11-2

Metabolic Pathways

Why pathways are broken down into little steps.

- Enzyme specificity allows only for breakage or formation of a few bonds at a time.
- Using pathways allows for multiple entry and exit points for metabolites.
- Pathways allow for finer control of energy input and output.
- Reactions are thermodynamically more efficient if carried out near equilibrium.

Chem 352, Lecture 7, Introduction to Metabolism 11

11-3

Metabolic Pathways

Why pathways are broken down into little steps.

- Enzyme specificity allows only for breakage or formation of a few bonds at a time.
- Using pathways allows for multiple entry and exit points for metabolites.
- Pathways allow for finer control of energy input and output.
- Reactions are thermodynamically more efficient if carried out near equilibrium.

Chem 352, Lecture 7, Introduction to Metabolism 11

11-4

Metabolic Pathways

Why pathways are broken down into little steps.

- Enzyme specificity allows only for breakage or formation of a few bonds at a time.
- Using pathways allows for multiple entry and exit points for metabolites.
- Pathways allow for finer control of energy input and output.
- Reactions are thermodynamically more efficient if carried out near equilibrium.

Chem 352, Lecture 7, Introduction to Metabolism 11

11-5

Metabolic Pathways

Pathways are regulated

- To control the flow of metabolites through a pathway

Chem 352, Lecture 7, Introduction to Metabolism 12

12-1

Metabolic Pathways

Pathways are regulated


- To control the flow of metabolites through a pathway

Chem 352, Lecture 7, Introduction to Metabolism 12

12-2

Metabolic Pathways

- Pathways are regulated
 - To control the flow of metabolites through a pathway




Chem 352, Lecture 7, Introduction to Metabolism 12

12-3

Metabolic Pathways

- Pathways are regulated
 - To control the flow of metabolites through a pathway




Chem 352, Lecture 7, Introduction to Metabolism 12

12-4

Metabolic Pathways

- Pathways are regulated
 - To control the flow of metabolites through a pathway



Chem 352, Lecture 7, Introduction to Metabolism 12

12-5

Metabolic Pathways


- Pathways are regulated
- Allosteric regulation responds to immediate conditions within the cell,
 - And have short term response times.
 - Reversible covalent modifications typically respond to extracellular signals,
 - And have longer term response times.

Chem 352, Lecture 7, Introduction to Metabolism 13

13-1

Metabolic Pathways

Pathways are regulated



Chem 352, Lecture 7, Introduction to Metabolism 13

13-2

Metabolic Pathways

Pathways are regulated

- Allosteric regulation responds to immediate conditions within the cell,
 - And have short term response times.
- Reversible covalent modifications typically respond to extracellular signals,
 - And have longer term response times.

Chem 352, Lecture 7, Introduction to Metabolism 13

13-3

Metabolic Pathways

Metabolic pathways represent the frontline for evolution.


- Genetic changes affect enzyme activities, which in turn, affect the flow of material through metabolic pathways.

Chem 352, Lecture 7, Introduction to Metabolism 14

14-1

Metabolic Pathways

Metabolic pathways represent the frontline for evolution.



Chem 352, Lecture 7, Introduction to Metabolism 14

14-2

Metabolic Pathways

Metabolic pathways represent the frontline for evolution.

- Genetic changes affect enzyme activities, which in turn, affect the flow of material through metabolic pathways.

Chem 352, Lecture 7, Introduction to Metabolism 14

14-3

Metabolic Pathways

Metabolic pathways represent the frontline for evolution.

- Genetic changes affect enzyme activities, which in turn, affect the flow of material through metabolic pathways.



Chem 352, Lecture 7, Introduction to Metabolism 14

14-4

Metabolic Pathways


Metabolic pathways represent the frontline for evolution.

- Genetic changes affect enzyme activities, which in turn, affect the flow of material through metabolic pathways.

Chem 352, Lecture 7, Introduction to Metabolism 14

14-5


Major Metabolic Pathways



Chem 352, Lecture 7, Introduction to Metabolism 15

15

Major Metabolic Pathways




Chem 352, Lecture 7, Introduction to Metabolism 16

16

Major Metabolic Pathways

- In many organisms, the various pathways are regulated through compartmentalization.



Chem 352, Lecture 7, Introduction to Metabolism 17

17

Spontaneity of Metabolic Reactions

- The spontaneity (favorability) of a chemical reaction can be determined from its Gibbs Free Energy (ΔG)

$$\Delta G = \Delta G^0 + RT \ln \left(\frac{[\text{products}]}{[\text{reactants}]} \right)$$

Chem 352, Lecture 7, Introduction to Metabolism 18

18

Spontaneity of Metabolic Reactions

Clicker Question:

A chemical reaction is spontaneous when the ΔG is

- A. Equal to 0
- B. Greater than 0
- C. Less than 0
- D. None of the above

Chem 352, Lecture 7, Introduction to Metabolism 19

19

Spontaneity of Metabolic Reactions

Under conditions of constant temperature and pressure there are two contributions to the free energy change

- Enthalpy, H
- Entropy, S

$$\Delta G = \Delta H - T\Delta S$$

Chem 352, Lecture 7, Introduction to Metabolism 20

20-1

Spontaneity of Metabolic Reactions

Under conditions of constant temperature and pressure there are two contributions to the free energy change

- Enthalpy, H
- Entropy, S

$$\Delta G = \Delta H - T\Delta S$$

Change in heat content
for the system

Chem 352, Lecture 7, Introduction to Metabolism 20

20-2

Spontaneity of Metabolic Reactions

Under conditions of constant temperature and pressure there are two contributions to the free energy change

- Enthalpy, H
- Entropy, S

$$\Delta G = \Delta H - T\Delta S$$

Change in heat content
for the system


Change in disorder
for the system

Chem 352, Lecture 7, Introduction to Metabolism 20

20-3

Spontaneity of Metabolic Reactions

The actual conditions within the cell must be considered when determining a ΔG value.



$$\Delta G = \Delta G^{\circ'} + RT \ln \left(\frac{[C][D]}{[A][B]} \right)$$

$Q = \frac{[C][D]}{[A][B]}$ is the mass action ratio

Chem 352, Lecture 7, Introduction to Metabolism 21

21

Spontaneity of Metabolic Reactions

- When $Q \approx K_{eq}$ a reaction is reversible.
- When $Q < K_{eq}$ a reaction is spontaneous and irreversible.
- When $Q > K_{eq}$ a reaction is non-spontaneous and irreversible

Chem 352, Lecture 7, Introduction to Metabolism 22

22

ATP


Adenosine Triphosphate (ATP) is one of the molecules used by a cell to store chemical energy.

- This energy is released by the hydrolysis of the two phosphate anhydride bonds.
- ATP is one of the ways that the energy released from catabolism is used to meet the energy requirements of anabolism

Chem 352, Lecture 7, Introduction to Metabolism 23

23-1

ATP



Chem 352, Lecture 7, Introduction to Metabolism 23

23-2

ATP

Adenosine 5'-triphosphate (ATP) is one of the molecules used by a cell to store chemical energy.

TABLE 10.1 Standard Gibbs free energies of hydrolysis for ATP, AMP, and pyrophosphate.

Reactants and products	ΔG° hydrolysis (kJ mol ⁻¹)
ATP + H ₂ O → ADP + P _i + H ⁺	-32
ATP + H ₂ O → AMP + PP _i + H ⁺	-45
AMP + H ₂ O → Adenosine + P _i	-13
PP _i + H ₂ O → 2 P _i	-29

P_i (inorganic phosphate) = HPO₄²⁻
 PP_i (pyrophosphate) = HP₂O₇³⁻

Inorganic phosphate (P_i) and Inorganic pyrophosphate (PP_i) are also shown.

Chem 352, Lecture 7, Introduction to Metabolism 23

23-3

ATP

Adenosine 5'-triphosphate (ATP) is one of the molecules used by a cell to store chemical energy.

TABLE 10.1 Standard Gibbs free energies of hydrolysis for ATP, AMP, and pyrophosphate.

Reactants and products	Hydrolysis of	ΔG° hydrolysis (kJ mol ⁻¹)
ATP + H ₂ O → ADP + P _i + H ⁺	Phosphate Anhydride	-32
ATP + H ₂ O → AMP + PP _i + H ⁺	Phosphate Anhydride	-45
AMP + H ₂ O → Adenosine + P _i	Phosphate Ester	-13
PP _i + H ₂ O → 2 P _i	Phosphate Anhydride	-29

P_i (inorganic phosphate) = HPO₄²⁻
 PP_i (pyrophosphate) = HP₂O₇³⁻

Inorganic phosphate (P_i) and Inorganic pyrophosphate (PP_i) are also shown.

Chem 352, Lecture 7, Introduction to Metabolism 23

23-4



23-5

- This energy is released by the hydrolysis of the two phosphate anhydride bonds.
- ATP is one of the ways that the energy released from catabolism is used to meet the energy requirements of anabolism

Chem 352, Lecture 7, Introduction to Metabolism 23

23-6



23-7



23-8

- This energy is released by the hydrolysis of the two phosphate anhydride bonds.
- ATP is one of the ways that the energy released from catabolism is used to meet the energy requirements of anabolism

Chem 352, Lecture 7: Introduction to Metabolism 21

23-9

ATP

Adenosine Triphosphate (ATP) is just one of the molecules used by a cell to store chemical energy.

- The other ribonucleotide triphosphates are also used for this same purpose.
 - Guanosine triphosphate (GTP)
 - Cytidine triphosphate (CTP)
 - Uridine triphosphate (UTP)

24-1

ATP

Question:

In a rat hepatocyte, the concentrations ATP, ADP and P_i are 3.4 mM, 1.3 mM and 4.8 mM, respectively. Calculate the Gibbs free energy for the hydrolysis of ATP in this cell. How does this compare to the standard free energy change?

24-2

ATP

Question:

In a rat hepatocyte, the concentrations ATP, ADP and P_i are 3.4 mM, 1.3 mM and 4.8 mM, respectively. Calculate the Gibbs free energy for the hydrolysis of ATP in this cell. How does this compare to the standard free energy change?

Reactants and products	$\Delta G'^{\circ}$ (kJ mol ⁻¹)
ATP + H ₂ O → ADP + P _i + H ⁺	-32
ATP + H ₂ O → ADP + PP _i + H ⁺	-45
ADP + H ₂ O → Adenosine + P _i	-13
PP _i + H ₂ O → 2 P _i	-29

P_i (inorganic phosphate) = 180 g mol⁻¹
 PP_i (pyrophosphate) = 180 g mol⁻¹

24-3

ATP


Adenosine Triphosphate (ATP) is just one of the molecules used by a cell to store chemical energy.

- The other ribonucleotide triphosphates are also used for this same purpose.
 - Guanosine triphosphate (GTP)
 - Cytidine triphosphate (CTP)
 - Uridine triphosphate (UTP)

24-4

ATP

The hydrolysis of ATP can be used to drive unfavorable reactions



25

ATP

•Phosphoryl-group-transfer potential

TABLE 10.3 Standard Gibbs free energies of hydrolysis for common metabolites

Metabolite	$\Delta G'^{\circ}$ hydrolysis (kJ mol ⁻¹)
Phosphoenolpyruvate	-62
1,3-Bisphosphoglycerate	-49
ATP to AMP + PP _i	-45
Phosphocreatine	-43
Phosphoguanine	-32
Acetyl CoA	-32
ATP to ADP + P _i	-32
Pyrophosphate	-29
Glucose 1-phosphate	-21
Glucose 6-phosphate	-14
Glycerol 3-phosphate	-9

Chem 352, Lecture 7, Introduction to Metabolism 26

26-1

ATP


•Phosphoryl-group-transfer potential

TABLE 10.3 Standard Gibbs free energies of hydrolysis for common metabolites

Metabolite	$\Delta G'^{\circ}$ hydrolysis (kJ mol ⁻¹)
Phosphoenolpyruvate	-62
1,3-Bisphosphoglycerate	-49
ATP to AMP + PP _i	-45
Phosphocreatine	-43
Phosphoguanine	-32
Acetyl CoA	-32
ATP to ADP + P _i	-32
Pyrophosphate	-29
Glucose 1-phosphate	-21
Glucose 6-phosphate	-14
Glycerol 3-phosphate	-9

Chem 352, Lecture 7, Introduction to Metabolism 26

26-2



26-3

ATP

•Phosphoryl-group-transfer potential

TABLE 10.3 Standard Gibbs free energies of hydrolysis for common metabolites

Metabolite	$\Delta G'^{\circ}$ hydrolysis (kJ mol ⁻¹)
Phosphoenolpyruvate	-62
1,3-Bisphosphoglycerate	-49
ATP to AMP + PP _i	-45
Phosphocreatine	-43
Phosphoguanine	-32
Acetyl CoA	-32
ATP to ADP + P _i	-32
Pyrophosphate	-29
Glucose 1-phosphate	-21
Glucose 6-phosphate	-14
Glycerol 3-phosphate	-9

Chem 352, Lecture 7, Introduction to Metabolism 26

26-4

ATP

•Nucleotidyl group transfer

• Used to activate substrates in ligase reactions

TABLE 10.3 Standard Gibbs free energies of hydrolysis for common metabolites

Metabolite	$\Delta G'^{\circ}$ hydrolysis (kJ mol ⁻¹)
Phosphoenolpyruvate	-62
1,3-Bisphosphoglycerate	-49
ATP to AMP + PP _i	-45
Phosphocreatine	-43
Phosphoguanine	-32
Acetyl CoA	-32
ATP to ADP + P _i	-32
Pyrophosphate	-29
Glucose 1-phosphate	-21
Glucose 6-phosphate	-14
Glycerol 3-phosphate	-9

Chem 352, Lecture 7, Introduction to Metabolism 27

27-1

ATP

- Nucleotidyl group transfer
- Used to activate substrates in ligase

Chem 352, Lecture 7, Introduction to Metabolism 27

27-2

ATP

- Nucleotidyl group transfer
- Used to activate substrates in ligase reactions

Chem 352, Lecture 7, Introduction to Metabolism 27

27-3

Thioesters as High Energy Compounds

- The thioester group also has a high energy for hydrolysis

Chem 352, Lecture 7, Introduction to Metabolism 28

28-1

Thioesters as High Energy Compounds

- The thioester group also has a high energy for hydrolysis

Metabolite	$\Delta G'^{\circ}$ hydrolysis (kJ mol ⁻¹)
Phosphoenolpyruvate	-62
1,3-Bisphosphoglycerate	-49
ATP to AMP + PP _i	-45
Phosphocreatine	-43
Phosphoguanine	-32
Acetyl CoA	-32
ATP to ADP + P _i	-32
Pyrophosphate	-29
Glucose 1-phosphate	-21
Glucose 6-phosphate	-14
Glycerol 3-phosphate	-9

Chem 352, Lecture 7, Introduction to Metabolism 28

28-2

Thioesters as High Energy Compounds

- The thioester group also has a high energy for hydrolysis


Metabolite	$\Delta G'^{\circ}$ hydrolysis (kJ mol ⁻¹)
Phosphoenolpyruvate	-62
1,3-Bisphosphoglycerate	-49
ATP to AMP + PP _i	-45
Phosphocreatine	-43
Phosphoguanine	-32
Acetyl CoA	-32
ATP to ADP + P _i	-32
Pyrophosphate	-29
Glucose 1-phosphate	-21
Glucose 6-phosphate	-14
Glycerol 3-phosphate	-9

Chem 352, Lecture 7, Introduction to Metabolism 28

28-3

Thioesters as High Energy Compounds

•The thioester group also has a high energy for hydrolysis




Chem 352, Lecture 7, Introduction to Metabolism 28

28-4

Thioesters as High Energy Compounds

•The thioester group also has a high energy for hydrolysis




Chem 352, Lecture 7, Introduction to Metabolism 28

28-5

Thioesters as High Energy Compounds

•The thioester group also has a high energy for hydrolysis



Chem 352, Lecture 7, Introduction to Metabolism 28

28-6

Reduced Coenzymes

Reduced coenzymes (NAD⁺, NADP, FAD, FMN, ubiquinone) provide another way to store chemical energy.


- They can be used to store the free energy that is released in oxidation reactions.
- The electrons released in these reactions are transferred to the coenzyme, usually in the form of a hydride (H⁻) ion.

Chem 352, Lecture 7, Introduction to Metabolism 29

29

Reduced Coenzymes

Reduction potentials can be used to measure the ability of a molecule to serve as a reducing agent in an oxidation/reduction reaction



Chem 352, Lecture 7, Introduction to Metabolism 30



30

Reduced Coenzymes

Reduction potentials can be measured with an electrochemical cell.

- The oxidation and reduction are separated by a wire.

The reduction of Cu^{2+} by Zn



Chem 352, Lecture 7, Introduction to Metabolism 31

31

Reduced Coenzymes

The change in the reduction potential for an oxidation/reduction reaction (ΔE°) can be used to determine the change in Free energy for the reaction.

$$\Delta E^{\circ} = E^{\circ}_{\text{electron acceptor}} - E^{\circ}_{\text{electron donor}}$$

$$\Delta G^{\circ} = -n\mathcal{F}\Delta E^{\circ}$$

n = number of electrons transferred

\mathcal{F} = Faraday's constant ($96,586 \text{ J V}^{-1} \text{ mol}^{-1}$)

Chem 352, Lecture 7, Introduction to Metabolism 32

32

Reduced Coenzymes

Standard reduction potentials, E° , are usually measured with respect to the reduction potential for of $2 \text{ H}^+(\text{aq}) \rightarrow \text{H}_2(\text{g})$

Reduction half-reaction	E° (V)
$\text{Acid/Cu} + \text{Cl}_2 + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{Acid} + 2\text{HCl}$	+0.68
$\text{Ferrodoxyn} + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{Ferredoxin} + 2\text{H}^+$	+0.43
$2\text{H}^+ + 2\text{e}^- \rightarrow \text{H}_2$ (at pH 7.0)	0.00
$\alpha\text{-Ketoglutarate} + \text{H}^+ + 2\text{e}^- + 2\text{H}^+ \rightarrow \text{Isocitrate}$	-0.38
$\text{Lipoic disulfide} + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{Lipoic dithiolane} (\text{LAD})$	-0.34
$\text{NADP}^+ + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{NADPH} + \text{H}^+$	-0.32
$\text{NAD}^+ + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{NADH} + \text{H}^+$	-0.32
$\text{Lipoic acid} + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{Lipoic dithiolane acid}$	-0.29
$\text{Ubiquinone (oxidized)} + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{Ubiquinol (reduced)}$	-0.10
$\text{FAD} + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{FADH}_2$	-0.21
$\text{FADH}_2 + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{FADH}_2$	-0.21
$\text{Acetate} + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{Acetate}$	-0.20
$\text{Pyruvate} + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{Pyruvate}$	-0.18
$\text{Malonate} + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{Malonate}$	-0.17

Chem 352, Lecture 7, Introduction to Metabolism 33

33-1

Reduced Coenzymes

Standard reduction potentials, E° , are usually measured with respect to the reduction potential for of $2 \text{ H}^+(\text{aq}) \rightarrow \text{H}_2(\text{g})$

Reduction half-reaction	E° (V)
$\text{Acid/Cu} + \text{Cl}_2 + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{Acid} + 2\text{HCl}$	+0.68
$\text{Ferrodoxyn} + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{Ferredoxin} + 2\text{H}^+$	+0.43
$2\text{H}^+ + 2\text{e}^- \rightarrow \text{H}_2$ (at pH 7.0)	0.00
$\alpha\text{-Ketoglutarate} + \text{H}^+ + 2\text{e}^- + 2\text{H}^+ \rightarrow \text{Isocitrate}$	-0.38
$\text{Lipoic disulfide} + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{Lipoic dithiolane} (\text{LAD})$	-0.34
$\text{NADP}^+ + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{NADPH} + \text{H}^+$	-0.32
$\text{NAD}^+ + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{NADH} + \text{H}^+$	-0.32
$\text{Lipoic acid} + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{Lipoic dithiolane acid}$	-0.29
$\text{Ubiquinone (oxidized)} + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{Ubiquinol (reduced)}$	-0.10
$\text{FAD} + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{FADH}_2$	-0.21
$\text{FADH}_2 + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{FADH}_2$	-0.21
$\text{Acetate} + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{Acetate}$	-0.20
$\text{Pyruvate} + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{Pyruvate}$	-0.18
$\text{Malonate} + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{Malonate}$	-0.17

Chem 352, Lecture 7, Introduction to Metabolism 33

33-2

Reduced Coenzymes

Standard reduction potentials, E° , are usually measured with respect to the reduction potential for of $2 \text{ H}^+(\text{aq}) \rightarrow \text{H}_2(\text{g})$

Reduction half-reaction	E° (V)
$\text{Acid/Cu} + \text{Cl}_2 + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{Acid} + 2\text{HCl}$	+0.68
$\text{Ferrodoxyn} + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{Ferredoxin} + 2\text{H}^+$	+0.43
$2\text{H}^+ + 2\text{e}^- \rightarrow \text{H}_2$ (at pH 7.0)	0.00
$\alpha\text{-Ketoglutarate} + \text{H}^+ + 2\text{e}^- + 2\text{H}^+ \rightarrow \text{Isocitrate}$	-0.38
$\text{Lipoic disulfide} + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{Lipoic dithiolane} (\text{LAD})$	-0.34
$\text{NADP}^+ + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{NADPH} + \text{H}^+$	-0.32
$\text{NAD}^+ + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{NADH} + \text{H}^+$	-0.32
$\text{Lipoic acid} + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{Lipoic dithiolane acid}$	-0.29
$\text{Ubiquinone (oxidized)} + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{Ubiquinol (reduced)}$	-0.10
$\text{FAD} + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{FADH}_2$	-0.21
$\text{FADH}_2 + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{FADH}_2$	-0.21
$\text{Acetate} + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{Acetate}$	-0.20
$\text{Pyruvate} + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{Pyruvate}$	-0.18
$\text{Malonate} + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{Malonate}$	-0.17

Chem 352, Lecture 7, Introduction to Metabolism 33

33-3

Reduced Coenzymes

Standard reduction potentials, E° , are usually measured with respect to the reduction potential for $2\text{H}^+(\text{aq}) \rightarrow \text{H}_2(\text{g})$

Reduction half-reaction	E° (V)
Acetyl-CoA + $\text{CoA-SH} + \text{H}^+ + 2\text{e}^- \rightarrow \text{Acetyl-CoA-SH} + \text{CoA}$	-0.8
Ferredoxin (cytochrome b_5) $\text{Fe}^{2+} + \text{e}^- \rightarrow \text{Fe}^{3+}$	-0.43
$2\text{H}^+ + 2\text{e}^- \rightarrow \text{H}_2(\text{g})$	0.00
α -Ketoglutarate + $\text{H}^+ + 2\text{e}^- \rightarrow \text{Succinate}$	-0.36
Lipoic lipoamide (FAD) + $2\text{H}^+ + 2\text{e}^- \rightarrow \text{Lipoic dihydrolipoamide (FADH}_2)$	-0.34
$\text{NAD}^+ + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{NADH} + \text{H}^+$	-0.32
$\text{NADP}^+ + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{NADPH} + \text{H}^+$	-0.32
Lactic acid + $2\text{H}^+ + 2\text{e}^- \rightarrow \text{Lactate}$	-0.29
Oxidation (reductase) $2\text{H}^+ + 2\text{e}^- \rightarrow \text{O}_2(\text{g})$ (reductase)	-0.22
FAD + $2\text{H}^+ + 2\text{e}^- \rightarrow \text{FADH}_2$	-0.22
FMN + $2\text{H}^+ + 2\text{e}^- \rightarrow \text{FMNH}_2$	-0.22
Acetate + $2\text{H}^+ + 2\text{e}^- \rightarrow \text{Ethanol}$	-0.20
Pyruvate + $2\text{H}^+ + 2\text{e}^- \rightarrow \text{Lactate}$	-0.18
Oxidation $2\text{H}^+ + 2\text{e}^- \rightarrow \text{H}_2(\text{g})$	-0.17

Chem 352, Lecture 7, Introduction to Metabolism 33

33-4

Reduced Coenzymes

Like ΔG , the observed change in the reduction potential for a reaction, (ΔE), can be determined relative to the change in the standard reduction potential, (ΔE°):

$$\Delta E = \Delta E^{\circ} - \frac{RT}{nF} \ln \left(\frac{[A_{ox}][B_{red}]}{[A_{red}][B_{ox}]} \right)$$


Chem 352, Lecture 7, Introduction to Metabolism 34

34-1

Reduced Coenzymes

Problem:

Determine the standard free energy change for the oxidation of $\text{NADH} + \text{H}^+$ by O_2 .




Chem 352, Lecture 7, Introduction to Metabolism 34

34-2

Reduced Coenzymes

Problem:

Determine the standard free energy change for the oxidation of $\text{NADH} + \text{H}^+$ by O_2 .



$$\Delta G^{\circ} = -n \Delta E^{\circ}$$

Chem 352, Lecture 7, Introduction to Metabolism 34

34-3

Reduced Coenzymes

TABLE 10.4

Standard reduction potentials of some important biological half-reactions

Reduction half-reaction	E° (V)
Cytochrome b_5 (microsome), $\text{Fe}^{2+} + \text{e}^- \rightarrow \text{Fe}^{3+}$	0.02
Formate + $2\text{H}^+ + 2\text{e}^- \rightarrow \text{Formate}$	0.03
Ubiquinone (Q) + $2\text{H}^+ + 2\text{e}^- \rightarrow \text{Ubiquinol (QH}_2)$	0.04
Cytochrome b (mitochondrial), $\text{Fe}^{2+} + \text{e}^- \rightarrow \text{Fe}^{3+}$	0.08
Cytochrome c_1 , $\text{Fe}^{2+} + \text{e}^- \rightarrow \text{Fe}^{3+}$	0.22
Cytochrome c , $\text{Fe}^{2+} + \text{e}^- \rightarrow \text{Fe}^{3+}$	0.23
Cytochrome a , $\text{Fe}^{2+} + \text{e}^- \rightarrow \text{Fe}^{3+}$	0.29
Cytochrome b , $\text{Fe}^{2+} + \text{e}^- \rightarrow \text{Fe}^{3+}$	0.36
Plastocyanin, $\text{Cu}^{2+} + \text{e}^- \rightarrow \text{Cu}^+$	0.37
$\text{NO}_3^- + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{NO}_2^- + \text{H}_2\text{O}$	0.42
Photosystem I (P700), $\text{P}_700 + \text{e}^- \rightarrow \text{P}_700^+$	0.43
$\text{I}_2/\text{O}_2 + 2\text{H}^+ + 2\text{e}^- \rightarrow \text{H}_2\text{O}$	0.77
Photosystem II (P680)	1.1

Chem 352, Lecture 7, Introduction to Metabolism 34

34-4

Reduced Coenzymes

TABLE 10.4 Standard reduction potentials of some important biological half-reactions

Reduction half-reaction	E'° (V)
Acetyl CoA + CO_2 + H^+ + $2e^- \rightarrow$ Pyruvate + CoA	-0.48
Ferredoxin (spinach), $\text{Fe}^{2+} + e^- \rightarrow \text{Fe}^{3+}$	-0.43
$2\text{H}^+ + 2e^- \rightarrow \text{H}_2$ (at pH 7.0)	-0.42
α -Ketoglutarate + CO_2 + $2\text{H}^+ + 2e^- \rightarrow$ Isocitrate	-0.38
Lipoyl dehydrogenase (FAD) + $2\text{H}^+ + 2e^- \rightarrow$ Lipoyl dehydrogenase (FADH ₂)	-0.34
NADPH + $2\text{H}^+ + 2e^- \rightarrow$ NADPH + H^+	-0.32
NADH + $2\text{H}^+ + 2e^- \rightarrow$ NADH + H^+	-0.32
Lipoic acid + $2\text{H}^+ + 2e^- \rightarrow$ Dihydrolipoic acid	-0.29
Glutathione (oxidized) + $2\text{H}^+ + 2e^- \rightarrow$ 2 Glutathione (reduced)	-0.23
FAD + $2\text{H}^+ + 2e^- \rightarrow$ FADH ₂	-0.22
FMN + $2\text{H}^+ + 2e^- \rightarrow$ FMNH ₂	-0.22
Acetaldehyde + $2\text{H}^+ + 2e^- \rightarrow$ Ethanol	-0.20
Pyruvate + $2\text{H}^+ + 2e^- \rightarrow$ Lactate	-0.18
Oxalosuccinate + $2\text{H}^+ + 2e^- \rightarrow$ Malate	-0.17

Chem 352, Lecture 7, Introduction to Metabolism 34

34-5

Reduced Coenzymes

Problem:
Determine the standard free energy change for the oxidation of NADH + H⁺ by O₂.


$$\frac{1}{2} \text{O}_2 + \text{NADH} + \text{H}^+ \rightarrow \text{H}_2\text{O} + \text{NAD}^+$$

$$\Delta G^{\circ} = -n \Delta E^{\circ}$$

Chem 352, Lecture 7, Introduction to Metabolism 34

34-6


Reduced Coenzymes



Chem 352, Lecture 7, Introduction to Metabolism 34

34-7


Reduced Coenzymes



Chem 352, Lecture 7, Introduction to Metabolism 34

34-8

Reduced Coenzymes




Chem 352, Lecture 7, Introduction to Metabolism 34

34-9

Reduced Coenzymes

Problem:

Determine the standard free energy change for the oxidation of NADH + H⁺ by O₂.



$$\Delta G^\circ = -n \cdot \Delta E^\circ$$

34-10

Reduced Coenzymes

Like ΔG , the observed change in the reduction potential for a reaction, (ΔE), can be determined relative to the change in the standard reduction potential, (ΔE°):


$$\Delta E = \Delta E^\circ - \frac{RT}{nF} \ln \left(\frac{[A_{ox}][B_{red}]}{[A_{red}][B_{ox}]} \right)$$

34-11

Reduced Coenzymes

Problem:

Determine the standard free energy change for the oxidation of succinate to fumarate by O₂.




35-1

Reduced Coenzymes

Problem:


Determine the standard free energy change for the oxidation of succinate to fumarate by O₂.



$$\Delta G^\circ = -n \cdot \Delta E^\circ$$

35-2

Reduced Coenzymes




35-3

Reduced Coenzymes

Problem:

Determine the standard free energy change for the oxidation of succinate to fumarate by O_2 .



Chem 352, Lecture 7, Introduction to Metabolism 35

35-4

Reduced Coenzymes

TABLE 10.4 Standard reduction potentials of some important biological half-reactions

Reduction half-reaction	E° (V)
Cytochrome b_5 (microsome), $Fe^{3+} + e^- \rightarrow Fe^{2+}$	0.02
Fumarate + $2H^+$ + $2e^- \rightarrow$ Succinate	0.03
Ubiquinone (Q) + $2H^+$ + $2e^- \rightarrow$ Ubiquinol	0.04
Cytochrome b (mitochondrial), $Fe^{3+} + e^- \rightarrow Fe^{2+}$	0.08
Cytochrome c_1 , $Fe^{3+} + e^- \rightarrow Fe^{2+}$	0.22
Cytochrome c , $Fe^{3+} + e^- \rightarrow Fe^{2+}$	0.23
Cytochrome f , $Fe^{3+} + e^- \rightarrow Fe^{2+}$	0.29
Cytochrome f , $Fe^{3+} + e^- \rightarrow Fe^{2+}$	0.36
Photosynthetic, $Cu^{2+} + e^- \rightarrow Cu^+$	0.37
$NO_3^- + 2H^+ + 2e^- \rightarrow NO_2^- + H_2O$	0.42
Photosystem I (P700), $Fe^{3+} + e^- \rightarrow Fe^{2+}$	0.43
$Fe^{3+} + e^- \rightarrow Fe^{2+}$	0.77
$1/2 O_2 + 2H^+ + 2e^- \rightarrow H_2O$	0.82
Photosystem II (P680)	1.1

Chem 352, Lecture 7, Introduction to Metabolism 35

35-5

Reduced Coenzymes

TABLE 10.4 Standard reduction potentials of some important biological half-reactions

Reduction half-reaction	E° (V)
Acetyl CoA + $CO_2 + H^+$ + $2e^- \rightarrow$ Pyruvate + CoA	-0.48
Ferredoxin (cytochrome), $Fe^{3+} + e^- \rightarrow Fe^{2+}$	-0.43
$2H^+ + 2e^- \rightarrow H_2$ (at pH 7.0)	-0.42
α -Ketoglutarate + $CO_2 + 2H^+ + 2e^- \rightarrow$ Isocitrate	-0.38
Lipoyl dehydrogenase (FAD) + $2H^+ + 2e^- \rightarrow$ Lipoyl dehydrogenase (FADH ₂)	-0.34
$NAD^+ + 2H^+ + 2e^- \rightarrow NADH + H^+$	-0.32
$NAD^+ + 2H^+ + 2e^- \rightarrow NADH + H^+$	-0.32
Lipoic acid + $2H^+ + 2e^- \rightarrow$ Dihydrolipoic acid	-0.29
Glutathione (oxidized) + $2H^+ + 2e^- \rightarrow$ 2 Glutathione (reduced)	-0.23
FAD + $2H^+ + 2e^- \rightarrow FADH_2$	-0.22
FMN + $2H^+ + 2e^- \rightarrow FMNH_2$	-0.22
Acetaldehyde + $2H^+ + 2e^- \rightarrow$ Ethanol	-0.20
Pyruvate + $2H^+ + 2e^- \rightarrow$ Lactate	-0.18
Oxalosuccinate + $2H^+ + 2e^- \rightarrow$ Malate	-0.17


Chem 352, Lecture 7, Introduction to Metabolism 35

35-6

Reduced Coenzymes

Problem:


Determine the standard free energy change for the oxidation of succinate to fumarate by O_2 .



Chem 352, Lecture 7, Introduction to Metabolism 35

35-7

Reduced Coenzymes



$$\Delta E = \Delta E^\circ - \frac{RT}{nF} \ln \left(\frac{[A_{ox}][B_{red}]}{[A_{red}][B_{ox}]} \right)$$


Chem 352, Lecture 7, Introduction to Metabolism 36

36

Next Up

Lecture 8 – Carbohydrate Metabolism

• Part I: Glycolysis



Chem 352, Lecture 7, Introduction to Metabolism 37
