


# Chem 352 – Lecture 8

## Carbohydrate Metabolism

### Part I: Glycolysis





# Introduction

Carbohydrate metabolism involves a collection of pathways.


- ♦ Glycolysis
  - Hexoses → 3-Carbon molecules
- ♦ Gluconeogenesis
  - 3-Carbon molecules → Hexoses
- ♦ Fermentation (anaerobic)
- ♦ Citric Acid Cycle (aerobic)
  - Oxidation all the way to  $\text{CO}_2 + \text{H}_2\text{O}$
- ♦ Pentose-Phosphate pathway
  - Hexose → Pentose



# Introduction

Carbohydrate metabolism involves a collection of pathways.

- ♦ Glycolysis
  - Hexoses → 3-Carbon molecules
- ♦ Gluconeogenesis
  - 3-Carbon molecules → Hexoses
- ♦ Fermentation (anaerobic)
- ♦ Citric Acid Cycle (aerobic)
  - Oxidation all the way to  $\text{CO}_2 + \text{H}_2\text{O}$
- ♦ Pentose-Phosphate pathway
  - Hexose → Pentose





# Introduction

Carbohydrate metabolism involves a collection of pathways.

- ♦ Glycolysis
  - Hexoses → 3-Carbon molecules
- ♦ Gluconeogenesis
  - 3-Carbon molecules → Hexoses
- ♦ Fermentation (anaerobic)
- ♦ Citric Acid Cycle (aerobic)
  - Oxidation all the way to  $\text{CO}_2 + \text{H}_2\text{O}$
- ♦ Pentose-Phosphate pathway
  - Hexose → Pentose


# The Glycolytic Reactions

- There are 10 reactions, which lead from glucose to pyruvate.
  - ✦ These reactions couple the lysis (splitting) and oxidation of hexose to the synthesis of 2 ATP's from ADP and  $P_i$ .





# The Glycolytic Reactions



• T  
fi



**Adenosine 5'-triphosphate (ATP<sup>-4</sup>)**




**Adenosine 5'-diphosphate (ADP<sup>-3</sup>)**




**Adenosine 5'-monophosphate (AMP<sup>-2</sup>)**

+



**Inorganic phosphate (P<sub>i</sub>)**

+



**Inorganic pyrophosphate (PP<sub>i</sub>)**


) of

2O



# The Glycolytic Reactions

• 7




**TABLE 10.1** Standard Gibbs free energies of hydrolysis for ATP, AMP, and pyrophosphate.


Reactants and products	$\Delta G^{\circ'}_{\text{hydrolysis}}$ (kJ mol <sup>-1</sup> )
ATP + H <sub>2</sub> O → ADP + P <sub>i</sub> + H <sup>+</sup>	-32
ATP + H <sub>2</sub> O → AMP + PP <sub>i</sub> + H <sup>+</sup>	-45
AMP + H <sub>2</sub> O → Adenosine + P <sub>i</sub>	-13
PP <sub>i</sub> + H <sub>2</sub> O → 2 P <sub>i</sub>	-29

P<sub>i</sub> (inorganic phosphate) = HPO<sub>4</sub><sup>2-</sup>

PP<sub>i</sub> (pyrophosphate) = HP<sub>2</sub>O<sub>7</sub><sup>3-</sup>




**Inorganic phosphate (P<sub>i</sub>)**





**Inorganic pyrophosphate (PP<sub>i</sub>)**

# The Glycolytic Reactions

• T  
fi




**Adenosine 5'-triphosphate (ATP<sup>-4</sup>)**





**Adenosine 5'-diphosphate (ADP<sup>-3</sup>)**

+




**Inorganic phosphate (Pi)**



**Adenosine 5'-monophosphate (AMP<sup>-2</sup>)**

+



**Inorganic pyrophosphate (PPi)**

)  
of

2O




# The Glycolytic Reactions

- There are 10 reactions, which lead from glucose to pyruvate.
  - ✦ These reactions couple the lysis (splitting) and oxidation of hexose to the synthesis of 2 ATP's from ADP and  $P_i$ .



# The Glycolytic Reactions

• There are 10 reactions which lead





# The Glycolytic Reactions

• There are 10 reactions which lead

**Oxidized form**

**Reduced form**

**TABLE 10.4** Standard reduction potentials of some important biological half-reactions

Reduction half-reaction	$E^{\circ'} \text{ (V)}$
Acetyl CoA + CO <sub>2</sub> + H <sup>+</sup> + 2e <sup>-</sup> → Pyruvate + CoA	-0.48
Ferredoxin (spinach), Fe <sup>3+</sup> + e <sup>-</sup> → Fe <sup>2+</sup>	-0.43
2 H <sup>+</sup> + 2e <sup>-</sup> → H <sub>2</sub> (at pH 7.0)	-0.42
α-Ketoglutarate + CO <sub>2</sub> + 2 H <sup>+</sup> + 2e <sup>-</sup> → Isocitrate	-0.38
Lipoyl dehydrogenase (FAD) + 2 H <sup>+</sup> + 2e <sup>-</sup> → Lipoyl dehydrogenase (FADH <sub>2</sub> )	-0.34
NADP <sup>+</sup> + 2 H <sup>+</sup> + 2e <sup>-</sup> → NADPH + H <sup>+</sup>	-0.32
NAD <sup>+</sup> + 2 H <sup>+</sup> + 2e <sup>-</sup> → NADH + H <sup>+</sup>	-0.32
Lipoic acid + 2 H <sup>+</sup> + 2e <sup>-</sup> → Dihydrolipoic acid	-0.29
Glutathione (oxidized) + 2 H <sup>+</sup> + 2e <sup>-</sup> → 2 Glutathione (reduced)	-0.23
FAD + 2 H <sup>+</sup> + 2e <sup>-</sup> → FADH <sub>2</sub>	-0.22
FMN + 2 H <sup>+</sup> + 2e <sup>-</sup> → FMNH <sub>2</sub>	-0.22
Acetaldehyde + 2 H <sup>+</sup> + 2e <sup>-</sup> → Ethanol	-0.20
Pyruvate + 2 H <sup>+</sup> + 2e <sup>-</sup> → Lactate	-0.18
Oxaloacetate + 2 H <sup>+</sup> + 2e <sup>-</sup> → Malate	-0.17

**TABLE 10.4** Standard reduction potentials of some important biological half-reactions


Reduction half-reaction	$E^{\circ'} \text{ (V)}$
Cytochrome <i>b</i> <sub>5</sub> (microsomal), Fe <sup>3+</sup> + e <sup>-</sup> → Fe <sup>2+</sup>	0.02
Fumarate + 2 H <sup>+</sup> + 2e <sup>-</sup> → Succinate	0.03
Ubiquinone (Q) + 2 H <sup>+</sup> + 2e <sup>-</sup> → QH <sub>2</sub>	0.04
Cytochrome <i>b</i> (mitochondrial), Fe <sup>3+</sup> + e <sup>-</sup> → Fe <sup>2+</sup>	0.08
Cytochrome <i>c</i> <sub>1</sub> , Fe <sup>3+</sup> + e <sup>-</sup> → Fe <sup>2+</sup>	0.22
Cytochrome <i>c</i> , Fe <sup>3+</sup> + e <sup>-</sup> → Fe <sup>2+</sup>	0.23
Cytochrome <i>a</i> , Fe <sup>3+</sup> + e <sup>-</sup> → Fe <sup>2+</sup>	0.29
Cytochrome <i>f</i> , Fe <sup>3+</sup> + e <sup>-</sup> → Fe <sup>2+</sup>	0.36
Plastocyanin, Cu <sup>2+</sup> + e <sup>-</sup> → Cu <sup>+</sup>	0.37
NO <sub>3</sub> <sup>-</sup> + 2 H <sup>+</sup> + 2e <sup>-</sup> → NO <sub>2</sub> <sup>-</sup> + H <sub>2</sub> O	0.42
Photosystem I (P700)	0.43
Fe <sup>3+</sup> + e <sup>-</sup> → Fe <sup>2+</sup>	0.77
1/2 O <sub>2</sub> + 2 H <sup>+</sup> + 2e <sup>-</sup> → H <sub>2</sub> O	0.82
Photosystem II (P680)	1.1

**NAD<sup>+</sup> (NADP<sup>+</sup>)**

**NADH (NADPH)**

# The Glycolytic Reactions

• There are 10 reactions which lead





# The Glycolytic Reactions

- There are 10 reactions, which lead from glucose to pyruvate.
  - ✦ These reactions couple the lysis (splitting) and oxidation of hexose to the synthesis of 2 ATP's from ADP and  $P_i$ .



# The Glycolytic Reactions

**TABLE 11.1** The reactions and enzymes of glycolysis

1. Glucose + ATP $\longrightarrow$ Glucose 6-phosphate + ADP + H <sup>+</sup>	Hexokinase, glucokinase
2. Glucose 6-phosphate $\rightleftharpoons$ Fructose 6-phosphate	Glucose-6-phosphate isomerase
3. Fructose 6-phosphate + ATP $\longrightarrow$ Fructose 1,6-bisphosphate + ADP + H <sup>+</sup>	Phosphofructokinase-1
4. Fructose 1,6-bisphosphate $\rightleftharpoons$ Dihydroxyacetone phosphate + Glyceraldehyde 3-phosphate	Aldolase
5. Dihydroxyacetone phosphate $\rightleftharpoons$ Glyceraldehyde 3-phosphate	Triose phosphate isomerase
6. Glyceraldehyde 3-phosphate + NAD <sup>+</sup> + P <sub>i</sub> $\rightleftharpoons$ 1,3-Bisphosphoglycerate + NADH + H <sup>+</sup>	Glyceraldehyde 3-phosphate dehydrogenase
7. 1,3-Bisphosphoglycerate + ADP $\rightleftharpoons$ 3-Phosphoglycerate + ATP	Phosphoglycerate kinase
8. 3-Phosphoglycerate $\rightleftharpoons$ 2-Phosphoglycerate	Phosphoglycerate mutase
9. 2-Phosphoglycerate $\rightleftharpoons$ Phosphoenolpyruvate + H <sub>2</sub> O	Enolase
10. Phosphoenolpyruvate + ADP + H <sup>+</sup> $\longrightarrow$ Pyruvate + ATP	Pyruvate kinase





# The Glycolytic Reactions

**TABLE 11.1** The reactions and enzymes of glycolysis

1. Glucose + ATP $\longrightarrow$ Glucose 6-phosphate + ADP + H <sup>+</sup>	Hexokinase, glucokinase
2. Glucose 6-phosphate $\rightleftharpoons$ Fructose 6-phosphate	Glucose-6-phosphate isomerase
3. Fructose 6-phosphate + ATP $\longrightarrow$ Fructose 1,6-bisphosphate + ADP + H <sup>+</sup>	Phosphofructokinase-1
4. Fructose 1,6-bisphosphate $\rightleftharpoons$ Dihydroxyacetone phosphate + Glyceraldehyde 3-phosphate	Aldolase
5. Dihydroxyacetone phosphate $\rightleftharpoons$ Glyceraldehyde 3-phosphate	Triose phosphate isomerase
6. Glyceraldehyde 3-phosphate + NAD <sup>+</sup> + P <sub>i</sub> $\rightleftharpoons$ 1,3-Bisphosphoglycerate + NADH + H <sup>+</sup>	Glyceraldehyde 3-phosphate dehydrogenase
7. 1,3-Bisphosphoglycerate + ADP $\rightleftharpoons$ 3-Phosphoglycerate + ATP	Phosphoglycerate kinase
8. 3-Phosphoglycerate $\rightleftharpoons$ 2-Phosphoglycerate	Phosphoglycerate mutase
9. 2-Phosphoglycerate $\rightleftharpoons$ Phosphoenolpyruvate + H <sub>2</sub> O	Enolase
10. Phosphoenolpyruvate + ADP + H <sup>+</sup> $\longrightarrow$ Pyruvate + ATP	Pyruvate kinase



# The Glycolytic Reactions

**TABLE 11.1** The reactions and enzymes of glycolysis


1. Glucose + ATP $\longrightarrow$ Glucose 6-phosphate + ADP + H <sup>+</sup>	Hexokinase, glucokinase
2. Glucose 6-phosphate $\rightleftharpoons$ Fructose 6-phosphate	Glucose-6-phosphate isomerase
3. Fructose 6-phosphate + ATP $\longrightarrow$ Fructose 1,6-bisphosphate + ADP + H <sup>+</sup>	Phosphofructokinase-1
4. Fructose 1,6-bisphosphate $\rightleftharpoons$ Dihydroxyacetone phosphate + Glyceraldehyde 3-phosphate	Aldolase
5. Dihydroxyacetone phosphate $\rightleftharpoons$ Glyceraldehyde 3-phosphate	Triose phosphate isomerase
6. Glyceraldehyde 3-phosphate + NAD <sup>+</sup> + P <sub>i</sub> $\rightleftharpoons$ 1,3-Bisphosphoglycerate + NADH + H <sup>+</sup>	Glyceraldehyde 3-phosphate dehydrogenase
7. 1,3-Bisphosphoglycerate + ADP $\rightleftharpoons$ 3-Phosphoglycerate + ATP	Phosphoglycerate kinase
8. 3-Phosphoglycerate $\rightleftharpoons$ 2-Phosphoglycerate	Phosphoglycerate mutase
9. 2-Phosphoglycerate $\rightleftharpoons$ Phosphoenolpyruvate + H <sub>2</sub> O	Enolase
10. Phosphoenolpyruvate + ADP + H <sup>+</sup> $\longrightarrow$ Pyruvate + ATP	Pyruvate kinase





# The Glycolytic Reactions

## • Reaction 1: Hexokinase



# The Glycolytic Reactions

## Clicker Questions:

There are four different hexokinase enzymes (I – IV) with differing  $K_M$  values:

Hexokinase	$K_M$
I, II, III	$10^{-4} - 10^{-6} \text{ M}$
IV	$10^{-2} \text{ M}$

Hexokinase IV, also known as glucokinase, is found in the liver. When the different tissues line up to take glucose from the blood, where is the liver in this lineup?

- A. First in line
- B. Last in line
- C. Somewhere in the middle



# The Glycolytic Reactions


## • Reaction 1: Hexokinase

- ✦ Different tissues have different **isoforms** of hexokinase.
  - The liver hexokinase, also called glucokinase, has the highest  $K_M$ , which reflects this organ's role in regulating blood glucose levels
- ✦ This reaction has a high negative  $\Delta G$ .
- ✦ Except for the liver, once phosphorylated, glucose cannot leave the cell.



# The Glycolytic Reactions

## • Reaction 1: Hexokinase



# The Glycolytic Reactions

Problem:

What is the standard free energy change,  $\Delta G^\circ$ , for the hexokinase reaction?



# The Glycolytic Reactions

## Problem:

What is the standard free energy change,  $\Delta G^\circ$ , for the hexokinase reaction?


**TABLE 10.3** Standard Gibbs free energies of hydrolysis for common metabolites

Metabolite	$\Delta G^\circ_{\text{hydrolysis}}$ (kJ mol <sup>-1</sup> )
Phosphoenolpyruvate	-62
1,3-Bisphosphoglycerate	-49
ATP to AMP + PP <sub>i</sub>	-45
Phosphocreatine	-43
Phosphoarginine	-32
Acetyl CoA	-32
ATP to ADP + P <sub>i</sub>	-32
Pyrophosphate	-29
Glucose 1-phosphate	-21
Glucose 6-phosphate	-14
Glycerol 3-phosphate	-9



# The Glycolytic Reactions


## • Reaction 1: Hexokinase





# The Glycolytic Reactions

## • Reaction 2: Glucose 6-Phosphate Isomerase





# The Glycolytic Reactions

## • Reaction 2: Glucose 6-Phosphate Isomerase

- ♦ In the cell, this reaction occurs near equilibrium



# The Glycolytic Reactions


## Clicker Questions:

If a reaction occurs near equilibrium, what does this say about the actual  $\Delta G$  for the reaction?

- A.  $\Delta G > 0$
- B.  $\Delta G \approx 0$
- C.  $\Delta G < 0$

# The Glycolytic Reactions


## • Reaction 3: Phosphofructokinase 1







# The Glycolytic Reactions


## • Reaction 3: Phosphofructokinase 1



Fructose 6-phosphate



Fructose 1,6-*bis*phosphate





# The Glycolytic Reactions


## • Reaction 3: Phosphofructokinase 1

- ✦ This enzyme catalyzes the first committed step in glycolysis.
- ✦ PFK-1 is regulated by numerous allosteric effectors.



# The Glycolytic Reactions


## • Reaction 4: Aldolase






# The Glycolytic Reactions

## • Reaction 4: Aldolase



# The Glycolytic Reactions


## • Reaction 4: Aldolase






# The Glycolytic Reactions

## • Reaction 4: Aldolase



# The Glycolytic Reactions


## • Reaction 4: Aldolase






# The Glycolytic Reactions

## • Reaction 4: Aldolase



# The Glycolytic Reactions

## •Reaction 5: Triose-Phosphate Isomerase





# The Glycolytic Reactions


## • Reaction 5: Triose-Phosphate Isomerase

- ✦ This reaction also occurs near equilibrium
- ✦ The reaction mechanism involves an endiol intermediate.

# Chemical Modes of Enzymatic Catalysis


- Acid/Base catalysis

- ✦ Triose phosphate isomerase illustrates both general acid and bases catalysis.






# Chemical Modes of Enzymatic Catalysis





is  
merase illustrates both  
ses catalysis.



# Chemical Modes of Enzymatic Catalysis


isomerase illustrates both  
 ses catalysis.






# The Glycolytic Reactions

## • Reaction 5: Triose-Phosphate Isomerase



# The Glycolytic Reactions


## •Reaction 5: Triose-Phosphate Isomerase





# The Glycolytic Reactions

## • Reaction 5: Triose-Phosphate Isomerase





# The Glycolytic Reactions

## • Reaction 5: Triose-Phosphate Isomerase


- ✦ This reaction also occurs near equilibrium
- ✦ The reaction mechanism involves an endiol intermediate.




# Chemical Modes of Enzymatic Catalysis

- Acid/Base catalysis


- ✦ Triose phosphate isomerase illustrates both general acid and bases catalysis.



# Chemical Modes of Enzymatic Catalysis





is  
merase illustrates both  
ses catalysis.






# Chemical Modes of Enzymatic Catalysis

isomerase illustrates both  
 ses catalysis.



# The Glycolytic Reactions

## •Reaction 6: Glyceraldehyde 3-Phosphate Dehydrogenase





# The Glycolytic Reactions


## • Reaction 6: Glyceraldehyde 3-Phosphate Dehydrogenase

- ✦ This reaction also takes place near equilibrium because the 1,3-bisphosphoglycerate is rapidly depleted.
- ✦  $\text{NAD}^+$  levels in the cell are typically low, so regeneration of  $\text{NAD}^+$ , is critical for this step in glycolysis.





# The Glycolytic Reactions

## • Reaction 7: Phosphoglycerate Kinase



**1,3-Bisphosphoglycerate**




**3-Phosphoglycerate**



# The Glycolytic Reactions

## Reaction 7: Phosphoenolpyruvate Kinase



# The Glycolytic Reactions

**TABLE 10.3** Standard Gibbs free energies of hydrolysis for common metabolites

Metabolite	$\Delta G^{\circ'}_{\text{hydrolysis}}$ (kJ mol <sup>-1</sup> )
Phosphoenolpyruvate	-62
1,3-Bisphosphoglycerate	-49
ATP to AMP + PP <sub>i</sub>	-45
Phosphocreatine	-43
Phosphoarginine	-32
Acetyl CoA	-32
ATP to ADP + P <sub>i</sub>	-32
Pyrophosphate	-29
Glucose 1-phosphate	-21
Glucose 6-phosphate	-14
Glycerol 3-phosphate	-9

Group-transfer potential (kJ mol<sup>-1</sup>)



Phospho

creatine


**High-energy compounds**

**Low-energy compounds**




# The Glycolytic Reactions

## Reaction 7: Phosphoenolpyruvate Kinase





# The Glycolytic Reactions

## • Reaction 7: Phosphoglycerate Kinase



**1,3-Bisphosphoglycerate**



**3-Phosphoglycerate**



# The Glycolytic Reactions


## • Reaction 7: Phosphoglycerate Kinase

- ♦ This enzyme is named for the reverse reaction.
- ♦ This reaction is an example of **substrate-level phosphorylation**.





# The Glycolytic Reactions

## •Reaction 8: Phosphoglycerate Mutase



**3-Phosphoglycerate**

**Phosphoglycerate  
mutase**




**2-Phosphoglycerate**




# The Glycolytic Reactions

## • Reaction 8: Phosphoglycerate Mutase



**3-Phosphoglycerate**

**Phosphoglycerate  
mutase**



**2-Phosphoglycerate**

# The Glycolytic Reactions

## • Reaction 8: Phosphoglycerate Mutase

### Clicker Questions:


What class of reaction is the phosphoglycerate mutase reaction?

- A. Transferase
- B. Lyase
- C. Isomerase
- D. Ligase
- E. Oxidoreductase




# The Glycolytic Reactions

## •Reaction 9: Enolase



# The Glycolytic Reactions


## •Reaction 10: Pyruvate Kinase





# The Glycolytic Reactions

## • Reaction 10: Pyruvate Kinase



# The Glycolytic Reactions

## • Reaction 10: Pyruvate Kinase

Group-transfer potential ( $\text{kJ mol}^{-1}$ )

70  
60  
50  
40  
30  
20  
10  
0

Phosph

**TABLE 10.3** Standard Gibbs free energies of hydrolysis for common metabolites

Metabolite	$\Delta G^{\circ'}_{\text{hydrolysis}}$ ( $\text{kJ mol}^{-1}$ )
Phosphoenolpyruvate	−62
1,3-Bisphosphoglycerate	−49
ATP to AMP + $\text{PP}_i$	−45
Phosphocreatine	−43
Phosphoarginine	−32
Acetyl CoA	−32
ATP to ADP + $\text{P}_i$	−32
Pyrophosphate	−29
Glucose 1-phosphate	−21
Glucose 6-phosphate	−14
Glycerol 3-phosphate	−9


High-energy compounds

Low-energy compounds




# The Glycolytic Reactions

## • Reaction 10: Pyruvate Kinase



# The Glycolytic Reactions



## •Reaction 10: Pyruvate Kinase





# The Glycolytic Reactions

## •Reaction 10: Pyruvate Kinase





# The Glycolytic Reactions


## • Reaction 10: Pyruvate Kinase

- ♦ Like phosphoglycerate kinase, this enzyme is named for the reverse reaction.
- ♦ This reaction is another example of **substrate-level phosphorylation**.



# The Fates of Pyruvate


- Pyruvate represents one of the major intersections in metabolism.





# The Fates of Pyruvate


- Pyruvate represents one of the major intersections in metabolism.





# The Fates of Pyruvate


- Fermentation is used to regenerate oxidized  $\text{NAD}^+$  when  $\text{O}_2$  cannot be utilized to do this.





# The Fates of Pyruvate



- Fermentation is used when  $\text{NAD}^+$  is not utilized to do this.





# The Fates of Pyruvate

- Fermentation is used when  $\text{NAD}^+$  is not available and is utilized to do this.





# Coenzymes and Vitamins (Chapter 7.7)

- Pyruvate decarboxylase uses the coenzyme thiamine pyrophosphate (TPP).

- ✦ thiamine pyrophosphate is synthesized from vitamin B<sub>1</sub> (thiamine)



**TABLE 7.2** Major coenzymes

Coenzyme	Vitamin source	Major metabolic roles	Mechanistic role
Adenosine triphosphate (ATP)	—	Transfer of phosphoryl or nucleotidyl groups	Cosubstrate
<i>S</i> -Adenosylmethionine	—	Transfer of methyl groups	Cosubstrate
Uridine diphosphate glucose	—	Transfer of glycosyl groups	Cosubstrate
Nicotinamide adenine dinucleotide (NAD <sup>+</sup> ) and nicotinamide adenine dinucleotide phosphate (NADP <sup>+</sup> )	Niacin	Oxidation-reduction reactions involving two-electron transfers	Cosubstrate
Flavin mononucleotide (FMN) and flavin adenine dinucleotide (FAD)	Riboflavin (B <sub>2</sub> )	Oxidation-reduction reactions involving one- and two-electron transfers	Prosthetic group
Coenzyme A (CoA)	Pantothenate (B <sub>3</sub> )	Transfer of acyl groups	Cosubstrate
Thiamine pyrophosphate (TPP)	Thiamine (B <sub>1</sub> )	Transfer of two-carbon fragments containing a carbonyl group	Prosthetic group
Pyridoxal phosphate (PLP)	Pyridoxine (B <sub>6</sub> )	Transfer of groups to and from amino acids	Prosthetic group
Biotin	Biotin	ATP-dependent carboxylation of substrates or carboxyl-group transfer between substrates	Prosthetic group
Tetrahydrofolate	Folate	Transfer of one-carbon substituents, especially formyl and hydroxymethyl groups; provides the methyl group for thymine in DNA	Cosubstrate
Adenosylcobalamin	Cobalamin (B <sub>12</sub> )	Intramolecular rearrangements	Prosthetic group
Methylcobalamin	Cobalamin (B <sub>12</sub> )	Transfer of methyl groups	Prosthetic group
Lipoamide	—	Oxidation of a hydroxyalkyl group from TPP and subsequent transfer as an acyl group	Prosthetic group
Retinal	Vitamin A	Vision	Prosthetic group
Vitamin K	Vitamin K	Carboxylation of some glutamate residues	Prosthetic group
Ubiquinone (Q)	—	Lipid-soluble electron carrier	Cosubstrate



# Coenzymes and Vitamins (Chapter 7.7)

- Pyruvate decarboxylase uses the coenzyme thiamine pyrophosphate (TPP).

- ✦ thiamine pyrophosphate is synthesized from vitamin B<sub>1</sub> (thiamine)





# Coenzymes and Vitamins (Chapter 7.7)

- Pyruvate decarboxylase uses the

coenzyme

(TPP)





# Coenzymes and Vitamins (Chapter 7.7)


- Pyruvate decarboxylase uses the coenzyme thiamine pyrophosphate (TPP).

- ✦ thiamine pyrophosphate is synthesized from vitamin B<sub>1</sub> (thiamine)



# Coenzymes and Vitamins (Chapter 7.7)

- Thiamine pyrophosphate is used in many decarboxylation reactions,
  - ✦ Including pyruvate decarboxylase






# Coenzyme

• Thiamine  
many

♦ Incl

er 7.7)


d in






# Coenzymes and Vitamins (Chapter 7.7)

- Thiamine pyrophosphate is used in many decarboxylation reactions,
  - ✦ Including pyruvate decarboxylase



# Free Energy Changes in Glycolysis


- The overall free energy change is negative.





# Free Energy Changes in Glycolysis

- The overall free energy change is negative.





# Regulation of Glycolysis


- The enzymes that catalyze the irreversible reactions are the primary sites of allosteric regulation.



# Regulation of Glycolysis


- The irreversible principle

ation.



# Regulation of Glycolysis

• The  
irre  
prin




ation.



# Regulation of Glycolysis


• The  
irre  
prin



ation.

# Regulation of Glycolysis


- The irre principle






# Regulation of Glycolysis

- The irre principle



# Regulation of Glycolysis


- The irre principle





# Regulation of Glycolysis

- The irreversible principle




Feedback inhibition  
"Citric Acid Cycle's needs  
for anabolic pathways  
are met"

# Regulation of Glycolysis

• The  
irre  
prin

ation.




In the Liver:  
"The hormone glucagon is not  
signaling a need to push  
glucose out into the blood."



# Regulation of Glycolysis

- The irreversible principle






# Regulation of Glycolysis

- In the liver, glucokinase, which is the liver's form of hexokinase, is also regulated by a regulatory protein that binds to glucokinase when fructose 6-phosphate levels are high.



# Regulation of Glycolysis

- In the livers that regulate blood glucose levels, fructose





# Regulation of Glycolysis

- In the liver, glucokinase, which is a liver form of hexokinase, is also regulated by a regulatory protein that binds to glucokinase when fructose 6-phosphate levels are high.



# Regulation of Glycolysis

- The liver is the only organ that is able to release glucose back into the blood


- ✦ It does so to satisfy the glucose need of other tissues, particularly the brain.



# Regulation of Glycolysis

- The liver is the only organ that is able to release glucose back into the blood

- ✦ It does so to satisfy the glucose need of





# Regulation of Glycolysis

- The liver is the only organ that is able to release glucose back into the blood

- ✦ It does so to satisfy the glucose need of other tissues, particularly the brain.



# Regulation of Glycolysis


The liver is the only organ that is able to release glucose back into the blood

- ✦ Under low blood glucose levels, the hormone glucagon signals the liver to halt glycolysis.
  - It does this using a signal transduction pathway.



# Transduction of Extracellular Signals

## •G-Proteins




# Regulation of Glycolysis

- The liver is the only organ that is able to release glucose back into the blood
  - ✦ Under low blood glucose levels, the hormone glucagon signals the liver to halt glycolysis.
    - It does this using a signal transduction pathway.



Reg


- The ability to block



is the hormone  
olysis.  
ion

# Regulation of Glycolysis

• The  
abl  
blo



In the Liver:  
"The hormone glucagon is not  
signaling a need to push  
glucose out into the blood."

It is  
to the


hormone  
colysis.  
ction



Regulation

• The ability to

block



is the

hormone

olysis.



# Regulation of Glycolysis

- The liver is the only organ that is able to release glucose back into the blood
  - ✦ Under low blood glucose levels, the hormone glucagon signals the liver to halt glycolysis.
    - It does this using a signal transduction pathway.



# Regulation of Glycolysis

- Regulation of Pyruvate Kinase


- ✦ Pyruvate Kinase is also regulated in response to glucagon.
  - Phosphorylation by Protein Kinase A in response to glucagon, lowers its activity.



# Regulation of Glycolysis

## • Regulation of Pyruvate Kinase


- ♦ Pyr
- resp
- Ph
- re





# Regulation of Glycolysis

• R






# Regulation of Glycolysis

## • Regulation of Pyruvate Kinase

- ♦ Pyr
- resp
- Ph
- re





# Regulation of Glycolysis

## Regulation of Pyruvate Kinase


- ♦ In addition to being allosterically inhibited by ATP, Pyruvate Kinase is also allosterically activated by fructose 6-phosphate.



# Regulation of Glycolysis

## Regulation of Pyruvate Kinase

- ♦ In liver, kidney, and red blood cells, pyruvate kinase is activated by ATP.






# Regulation of Glycolysis

## Regulation of Pyruvate Kinase

- ♦ In the presence of ATP, pyruvate kinase activity is inhibited

Pyruvate kinase activity



regulated by


pyruvate]



# Regulation of Glycolysis

## Regulation of Pyruvate Kinase

- ♦ In liver, kidney, and red blood cells, pyruvate kinase is activated by ATP.





# Other Points

## Skip Section 11.6

- ♦ Entry of other sugars into glycolysis

## Skip Section 11.7

- ♦ Entner-Doudoroff Pathway in Bacteria, which lack PFK-1

# Review of Glycolysis

## Questions:

What is the metabolic purpose behind the glycolytic pathway?



# Review of Glycolysis

## Questions:

The “-lysis” part of glycolysis means “to split”. Using structural formulas, draw the chemical equation for the reaction in which a 6-carbon molecule is split into two 3-carbon molecules.

# Review of Glycolysis

## Questions:

Even though the glycolytic pathway can be used in the absence of oxygen, there is one reaction in which and oxidation occurs.

- Using structural formulas, draw the chemical equation for this reaction.
- What oxidizing reagent is used in this reaction?



# Review of Glycolysis

## Questions:

When oxygen is available and can be used, the oxidizing agent using in the previously described reaction is reoxidized by the electron transport chain.

- Using structural formulas, show the reaction or combination or reactions that is used to reoxidize this agent when oxygen cannot be utilized.
- What is the name of the pathway you drew?

# Next Up

## •Lecture 8 - Carbohydrate Metabolism

- ♦ Part II: Gluconeogenesis, Pentose Phosphate Pathway, and Glycogen Metabolism (Moran et al., Chapter 12)