Name	

Chem 352 - Spring 2011 - Exam IV

In	photosynthesis,
a.	Light energy is used to excite electrons in the photoreaction systems and remove them from a pair of chlorophyll molecules called the
b.	For purple bacterial, the energetically excited electrons from Photosystem II are used to reduce a mobile electron carrier named
c.	For green sulfur bacteria, the excited electrons from Photosystem I are used to reduce one of either two mobile carriers, or
d.	For both the purple bacteria and the green sulfur bacteria, describe, in one or two sentences, how the absorption of light energy is coupled to the synthesis ATP from ADP and P_i .
e.	What two products from the light reactions of photosynthesis in plants are used to fix CO ₂ in the <i>Calvin Cycle</i> ? and
f.	Pick an example of a reaction from the Calvin Cycle that illustrates the use of one of these two products and another reaction that does not. Use structural formulas to write a balanced chemical equation for your two chosen reactions. Name the intermediates and the enzymes involved in each.
	i.
	ii.
g.	What other metabolic pathway shares the same reactions found in the reduction phase of the <i>Calvin cycle</i> ?

	. 11.
2. Fatty acids are synthesized by adding two carbon units to a	SIOWING ACVICINATION.

a. *Using structural formulas*, show the reactions involved in elongating an acyl chain by two carbon atoms.

b. Fatty acid synthesis shares a number of similar features to β -oxidation (the degradation of fatty acids), but in reverse. Describe two features of fatty acid biosynthesis that distinguish it from the reverse of β -oxidation.

i.

ii.

c. Where in a eukaryotic cell does fatty acid synthesis occur?

d. When eukaryotes convert carbohydrates to fatty acids, the glycolytic pathway is used to convert glucose to pyruvate, which is then converted to acetyl-CoA using pyruvate dehydrogenase. Describe where in the cell this acetyl-CoA is produced and indicate how it is transported to the location where fatty acid biosynthesis is taking place.

3.	ma util	humans, the acetyl-CoA produced from the β -oxidation of fatty acid cannot be used as starting terial for the synthesis of glucose, however, under conditions of starvation, acetyl-CoA can be lized by the liver to produce a group of molecules that circulate to the tissues and are used as an ernative to glucose as an fuel source.	
	a.	What is the name for this group of molecules?	
	b.	Draw the structural formulas for, and name two examples of molecules that are members of this group.	
	c.	The reactions in the pathway that leads to the synthesis of these molecules are also found in another important biosynthetic pathway. Name this pathway this?	
4.		me the <i>glycolytic</i> or <i>citric acid cycle</i> intermediate that is the starting point for the biosynthesis of the following amino acids:	
		asparagine	
		alanine	
		aspartate	
		glycine	
		serine	
		proline	
		glutamate	
	a.	Pick two of these, one from glycolysis and one from the citric acid cycle, and <i>using structural formulas</i> , draw out the pathway leading to their biosynthesis.	
		i.	

ii.

5. Both adenosine monophosphate and guanosine monophosphate are synthesized from inosine monophosphate. Draw the structure of adenosine monophosphate and indicate the source of each of the atoms in ring of the nucleotide base. 6. Terrestrial animals dispose of their excess nitrogen as urea. a. Draw the structural formula for urea b. Urea is made by the urea cycle. *Using structural formulas*, draw the reactions for the urea cycle: c. Two of the intermediates in the urea cycle are α -amino acids that are not members of the 20 used to make proteins. Name these: i. ii.

Extra Credit:

1. Ask the one question that you wanted me to ask, but I did not ask. (Up to 2 points will be awarded for an insightful, probing and well-worded question.)

2. Answer the question you posed in part 1. (Up to 1 point will be awarded for answering your question correctly.)